

**RESEARCHES REGARDING THE INFLUENCE OF
FORERUNNER PLANT x FERTILISATION LEVEL INTERACTION
ON BIOMASS ACCUMULATION IN WINTER WHEAT
CULTIVATED ON BROWN LUVIC SOILS**

Ardelean Ileana, Bandici Gheorghe-Emil*

*University of Oradea, Faculty of Environmental Protection, 26 Gen. Magheru St., 410048 Oradea;
Romania, e-mail: gbandici@yahoo.com

Abstract

The forerunner plant is a decisive factor influencing growth and development of wheat.

Plant growth is fundamental in obtaining yield and is related vegetation and technological factors, the level of yield being reflected in the intensity of phytomass accumulation.

In the majority of cases, total growth of green mass is considered on the assumption that a maximum yield is obtained by increasing total dry weight biomass production and by a favourable repartition of it among plant's organs.

Key words: crop rotation, fertilization, phenophase, phytomass, forerunner plant, climatic conditions

INTRODUCTION

Crop rotation together with other appropriate agricultural practices contribute to the favourableness of growth and development conditions of wheat root system, to an improved synthesis of specific organic compounds and their improved translocation to plant's organs (*Lazany J., 2000; Bandici Gh., 1997, 2001*) Finally, all the enumerated conditions lead to improved efficiency per area unit.

Zamfirescu, (1977) stressed out that the accumulation process of total green mass relies on the absorption of nutrients from soil through plant's root system. In turn, root system depends on the level of foliar system functioning conditioned by water and nutrients' provision of soil. (*Lazany J., 2003*).

The importance of rationale fertilization and of the forerunner plant on growths and development of plants is stated by many authors: *Popescu Ana (1980), Zăhan P. și Zăhan Rodica (1989b)*.

Most of the reserches were centred on the influence of crop rotation on the yields, namely on the phytomass accumulation. the crop rotations with regard to wheat was very satisfactory in this order as forerunner plant: pea, beans, winter rape, bots, linseed, soja, red clover, potato, sugar beet, sunflower, corn etc. (*Munteanu L., and all., 2008*).

Bîlteanu Gh., (1993) after long tests demonstrated the importance of crop rotation on wheat yields on brown-red soils in Romanian Plain. On

clay-illuvial podzols, the introduction of ameliorative plants such as red clover represented an element of utmost importance for increase of the wheat yield.

Dincă, D., (1982), made some references on the role of crop rotation on wheat yield and on the organic accumulation in whole plant and grains.

It is demonstrated that after 10-year monoculture, wheat yield decreases continuously in comparative with rotations. it fluctuates as a consequence of changing climatic conditions. under such circumstances, fertilization does not induce a significant yield increase. a particularly important problem is linked to wheat crop increment, which must fit the rising consumption needs of world population (*Zăhan P. și Zăhan Rodica, 1989a*).

MATERIAL AND METHODS

The research was set at Agrozootechnical Researches Experimental Station (A.R.E.S) Oradea (Romania), between 2009 - 2010. This period was characterized by an alternation of favourable climatic conditions during 2009 (a normal climatic year) and dry climatic conditions during 2010, less favorable, consequently. The experimental design was polyfactorial in subdivides stands using as factors interaction: climatic conditions versus fertilization level. The influence of later factors was studied on the dynamics of winter wheat accumulation cultivated on brown luvic soils. As biological material, the Delia race of wheat was employed.

Experimental results (biomass accumulation) were analyzed by ANOVA (analysis of variance) and expressed as g of dry weight/10 plants (Bandici, 2007).

RESULTS AND DISCUSSIONS

The present paper presents the correlation between biomass accumulation and forerunner plant quality in winter wheat cultivation, function of climatic conditions current year.

Table 1 presents the biomass accumulation during the less favourable year 2009. The results show that the forerunner plant do not influence biomass accumulation increase as winter wheat advances in vegetation. The increase is dependent on the properties and quality of the forerunner plant, the best yields being obtained after the cultivation of pea. Thus, in wheat monoculture the biomass accumulation increases from 0.96 g. dry weight/10 plants (at beginning winter) to 45.40 g. dry weight/10 plants at complete ripening. If corn is used as forerunner plant, but specially after pea, the increases in yield are superior, varying between 0.87-60.43 g.

dry weight/10 plants after corn and between 1.50-66.97 g. dry weight/10 plants after pea, respectively as winter wheats' forerunner plants.

The phenomenon keeps same pace but at values in 2010, a less favourable year (table 2). The biomass accumulation varies in wheat monoculture between 0.20-35.60 g. dry weight/10 plants and 0.23-42.83 g. dry weight/10 plants after corn and between 0.20-47.10 g. dry weight/10 plants after pea. Regardless to climatic conditions, at a different scale, a positive correlation was found between phenophase x fertilization level: as wheat advances to maturity, there is a progressive accumulation of total biomass in seeds (table 3 and 4).

Concerning 2009, a less favorable year, results show a very significant increase in biomass at the beginning of winter in unfertilized alternative (1.00 g dry weight/10 plants). At the complete ripening there is an increase to 49.92 g dry weight/10 plants, in unfertilized alternative (*Zăhan, Rodica Zăhan, 1989*).

In 2010, a less favorable year, biomass accumulation decreases as compared to a favorable year, taking values in a narrower, range between 0.22 g dry weight/10 plants at beginning of winter and 31.32 g dry weight/10 plants at compared ripening in unfertilized alternative. It is worth to mention that in 2009 (less favorable year) the quantity of accumulated biomass was directly proportional with fertilization on level as it was rising during study period (49.92 g dry weight/10 plants) in unfertilized alternative and 69.49 g dry weight/10 plants in fertilized alternative, organo-mineral complex was used in all experimental alternative as fertilized.

In a less favorable year, 2010, biomass accumulation decreased depending on fertilization level (31.32 g dry weight/10 plants in unfertilized alternative and 51.80 g dry weight/10 plants in fertilized alternative using the same organo-mineral complex as fertilizer).

It is worth to mention that during the two research years, a negative correlation was found between percent participation of compound synthesized before fructification and seeds formation and fertilization level.

As fertilization level increases, the percent participation of compounds synthesized before fructification insides formation decreases. This phenomen is more accentuated in unfertilized alternatives as compared with fertilized with organo-mineral complex and is influenced unfavorable climatic conditions.

Table 1

The effect of the forerunner plant x phenophase on winter wheat biomass accumulation dynamics, on brown luvisc soils Oradea (Romania) 2009

Phenophase	Total dry weight biomass, seeds and straw g./10 plants											
	Forerunner plant											
	Wheat			Corn			Pea III			Pea IV		
	Total s d.W.	Seeds	Straw	Total s d.W.	Seeds	Straw	Total s d.W.	Seeds	Straw	Total s d.W.	Seeds	Straw
At winter beginning	0.96	-	0.96	0.87	-	0.87	0.87	-	0.87	1.50	-	1.50
At the end of winter	1.07	-	1.07	1.07	-	1.07	1.57	-	1.57	2.20	-	2.20
The beginning of vegetation	1.93	-	1.93	2.96	-	2.96	4.27	-	4.27	4.07	-	4.07
The formation of first interned	5.27	-	5.27	5.10	-	5.10	7.80	-	7.80	6.27	-	6.27
Straw elongation	13.47	-	13.47	19.17	-	19.17	17.17	-	17.17	14.00	-	14.00
The formation of spike	31.50	-	31.50	29.50	-	29.50	54.60	-	54.60	49.53	-	49.53
Beginning of seeds formation	34.63	-	34.63	42.83	-	42.83	57.90	-	57.90	54.63	-	54.63
Early ripening	39.67	10.00	29.67	52.90	13.33	39.57	59.93	14.33	45.60	56.80	11.00	45.80
Incomplete ripening	45.33	12.53	32.80	57.10	15.93	41.17	62.50	21.67	40.83	62.67	18.67	44.00
Complete ripening	45.40	15.67	29.73	60.43	24.33	36.10	63.60	27.00	36.60	66.97	30.67	36.30
LSD 5%				0.071 g/10 plants d.w.						2.7 g/10 plants d.w.		
LSD 1 %	For total plant biomass			0.093 g/10 plants d.w.			For seeds:			3.6 g/10 plants d.w.		
LSD 0,1 %				0.119 g/10 plants d.w.						4.7 g/10 plants d.w.		

- % participation of synthesised compounds prior to fructification and seeds' formation: Wheat = 14.1%; Pea III = 36.8%; Corn = 15.7%; Pea IV = 33.5%;

Statistical significations: for *Total plant biomass*: under 0.071 = insignificant (-); 0.071-0.093 = significant (*); 0.093 - 0.119 = distinct significant (**); over 0.119 = very significant (***)

- for *seeds*: under 2.7 = insignificant (-); 2.7-3.6 = significant (*); 3.6 - 4.7 = distinct significant (**); over 4.7 = very significant (***)

Table 2.

The effect of the forerunner plant x phenophase on winter wheat biomass accumulation dynamics. on brown luvisc soils Oradea (Romania) 2010

Phenophase	Total dry weight biomass. seeds and straw g./10 plants											
	Forerunner plant											
	Wheat			Corn			Pea III			Pea IV		
	Total s d.W.	Seeds	Straw	Total s d.W.	Seeds	Straw	Total s d.W.	Seeds	Straw	Total s d.W.	Seeds	Straw
At winter beginning	0.20	-	0.20	0.23	-	0.23	0.30	-	0.30	0.20	-	0.20
At the end of winter	0.37	-	0.37	0.37	-	0.37	0.40	-	0.40	0.37	-	0.37
The beginning of vegetation	0.47	-	0.47	0.47	-	0.47	0.50	-	0.50	0.47	-	0.47
The formation of first interned	0.80	-	0.80	0.97	-	0.97	1.33	-	1.33	0.93	-	0.93
Straw elongation	3.73	-	3.73	3.43	-	3.43	4.37	-	4.37	4.33	-	4.33
The formation of spike	8.60	-	8.60	9.47	-	9.47	11.0	-	11.00	10.60	-	10.60
Beginning of seeds formation	24.83	-	24.83	25.50	-	25.50	34.03	-	34.03	33.13	-	33.13
Early ripening	33.87	9.83	24.04	36.00	8.50	27.50	35.73	11.33	24.40	41.33	10.83	30.50
Incomplete ripening	35.37	11.70	23.67	41.87	11.00	30.87	40.73	13.90	26.83	45.33	12.17	33.16
Complete ripening	35.60	13.37	22.23	42.83	12.80	30.03	43.43	17.50	25.93	47.10	15.97	31.13
LSD 5%	0.073 g/10 plants d.w.						2.7 g/10 plants d.w.					
LSD 1 %	.For total plant biomass: 0.096 g/10 plants d.w.						For seeds: 3.6 g/10 plants d.w.					
LSD 0.1 %	0.123 g/10 plants d.w.						4.7 g/10 plants d.w.					

- % participation of synthesised compounds prior to fructification and seeds' formation: Wheat = 10.5%; Pea III = 28.3%;
Corn = 17.4%; Pea IV = 8.0 %.

Statistical significations:

- for *Total plant biomass*: under 0.073 = insignificant (-); 0.073-0.096 = significant (*); 0.096-0.123 = distinct significant (**); over 0.123 = very significant (***)

- for *seeds*: under 2.7 = insignificant (-); 2.7-3.6 = significant (*); 3.6-4.7 = distinct significant (**); over 4.7 = very significant (***)

Table 3

The effect of fertilisation level x phenophase interaction on winter wheat dry weight biomass accumulation dynamics on brown luvic soils.
Oradea 2009

Phenophase	Total dry weight biomass, seeds and straw g./10 plants								
	Fertilisation level								
	N ₀ P ₀			N ₁₂₀ P ₈₀			N ₁₀₀ P ₈₀ + 10t/ha manure		
	Total s d.W.	Seeds	Straw	Total s d.W.	Seeds	Straw	Total s d.W.	Seeds	Straw
At winter beginning	1.00	-	1.00	1.17	-	1.17	0.97	-	0.97
At the end of winter	1.22	-	1.22	1.55	-	1.55	1.65	-	1.65
The beginning of vegetation	2.65	-	2.65	2.62	-	2.62	4.65	-	4.65
The formation of first interned	4.77	-	4.77	6.10	-	6.10	7.45	-	7.45
Straw elongation	11.62	-	11.62	16.85	-	16.85	19.37	-	19.37
The formation of spike	36.70	-	36.70	43.05	-	43.05	44.10	-	44.10
Beginning of seeds formation	43.62	-	43.62	49.55	-	49.55	49.31	-	49.31
Early ripening	47.62	9.50	38.12	52.90	13.75	39.15	56.45	13.25	43.20
Incomplete ripening	49.82	15.75	34.07	57.32	18.45	38.87	63.55	17.40	46.15
Complete ripening	49.92	21.00	28.92	57.95	24.50	33.45	69.49	27.75	41.74
DL 5%	0.063 g/10 plants d.w.						2.3 g/10 plants d.w.		
DL 1%	For total plant biomass: 0.083 g/10 plants d.w.						For seeds: 3.1 g/10 plants d.w.		
DL 0.1%	0.106 g/10 plants d.w.						4.0 g/10 plants d.w.		

- % participation of synthesised compounds prior to fructification and seeds' formation: N₀P₀ = 33.7 %; N₁₂₀P₈₀ = 32.5 %;
N₁₀₀P₈₀ + 10t/ha manure = 15.3 %

Statistical significations:

- for *Total plant biomass*: under 0.063 = insignificant (-); 0.063-0.083 = significant (*); 0.083-0.106 = distinct significant (**); over 0.106 = very significant (***)

- for *seeds*: under 2.3 = insignificant (-); 2.3-3.1 = significant (*); 3.1-4.0 = distinct significant (**); over 4.0 = very significant (***)

Table 4

The effect of fertilisation level x phenophase interaction on winter wheat dry weight biomass accumulation dynamics on brown luvisols.
Oradea 2010

Phenophase	Total dry weight biomass. seeds and straw g./10 plants								
	Fertilisation level								
	N ₀ P ₀			N ₁₂₀ P ₈₀			N ₁₀₀ P ₈₀ + 10t/ha manure		
	Total s d.W.	Seeds	Straw	Total s d.W.	Seeds	Straw	Total s d.W.	Seeds	Straw
At winter beginning	0.22	-	0.22	0.22	-	0.22	0.25	-	0.25
At the end of winter	0.35	-	0.35	0.37	-	0.37	0.40	-	0.40
The beginning of vegetation	0.45	-	0.45	0.47	-	0.47	0.50	-	0.50
The formation of first interned	0.90	-	0.90	0.95	-	0.95	1.17	-	1.17
Straw elongation	2.45	-	2.45	4.47	-	4.47	4.97	-	4.97
The formation of spike	5.82	-	5.82	11.30	-	11.30	12.62	-	12.62
Beginning of seeds formation	23.00	-	23.00	30.07	-	30.07	35.05	-	35.05
Early ripening	26.45	6.03	20.42	38.97	11.85	27.12	44.77	12.50	32.27
Incomplete ripening	30.22	7.35	22.87	43.05	14.07	28.98	49.20	15.15	34.05
Complete ripening	31.32	9.72	21.60	43.60	15.70	27.90	51.80	19.30	32.50
DL 5%				0.063 g/10 plants d.w.			2.3 g/10 plants d.w.		
DL 1 %				0.083 g/10 plants d.w.			For seeds: 3.1 g/10 plants d.w.		
DL 0.1 %				0.106 g/10 plants d.w.			4.0 g/10 plants d.w.		

- % participation of synthesised compounds prior to fructification and seeds' formation: N₀P₀ = 11.2 %; N₁₂₀P₈₀ = 9.8 %;
N₁₀₀P₈₀ + 10t/ha manure = 7.9 %

Statistical significations:

- for *Total plant biomass*: under 0.063 = insignificant (-); 0.063-0.083 = significant (*); 0.083-0.106 = distinct significant (**); over 0.106 = very significant (***)

- for *seeds*: under 2.3 = insignificant (-); 2.3-3.1 = significant (*); 3.1-4.0 = distinct significant (**); over 4.0 = very significant (***)

Thus, in 2009, a favorable year, increasing the fertilization had as result a decrease of participation percentage from 33.7 g dry weight/10 plants in unfertilized alternative to 15.3 g dry weight/10 plants in fertilized with organo-mineral complex alternatives.

In 2010, considered an unfavorable year, the decreases varied between 11.2 g dry weight/10 plants in unfertilized and 7.9 g dry weight/10 plants in fertilized alternatives using organo-mineral complex of fertilizers.

CONCLUSIONS

The forerunner plants' quality has a positive effect on total biomass accumulation as compared with wheat monoculture, the obtained values being conditioned by a higher favourableness of climatic factors.

With regard to biomass accumulation in seeds, it was positively influenced by the quality of the forerunner plant, being higher in correlation with the forerunner plants' better qualities.

REFERENCES

1. Bandici, G., E., 1997, *Contribuții la stabilirea influenței premergătoarei și a fertilizării asupra dinamicii acumulării biomasei, la grâul de toamnă, cultivat pe soluri cu exces temporar de umiditate, în centrul Câmpiei de Vest a României*. Doctoral thesis. University of Agriculture Sciences and Veterinary Medicine Cluj-Napoca, Romania [in Romanian], p.158.
2. Bandici, G., E., P., Guș, 2001, *Dinamica acumulării de biomasă la grâul de toamnă*. University of Oradea Press, p.107.
3. Bîlteanu, G., 1993, *Fitotehnie*, Ceres Printing House. Bucharest, p.457.
4. Dincă, D., 1982, *Asolamentele agriculturii moderne*. Ceres Printing House, Bucharest, p.257.
5. Lazany, J., 2000, *Soil fertility management in Westik's crop rotation experiment. Role of fertilizers in Sustainable Agriculture*. CIEC Conference. pp.77-80, pp.255.
6. Lazany, J., 2003, *Differences in soil carbon content in the treatments of Westik's crop rotation experiment. Natural resources and sustainable development*. International scientific session and reviewed papers. Oradea-Debrecen, pp.119-120, pp.288.
7. Muntean, L.S., Cernea, S., Morar, G., Duda, M., Vârban, D., Muntean, S., 2008: *Fitotehnie*. Academic Pres Printing House, Cluj-Napoca, pp. 83-135.
8. Popescu Ana - 1980 - "*Procesul de fixare biologică a N atmosferic și factorii care îl condiționează*". Probleme de Agrofit. teor. și practică, nr.1, vol. II, pp. 35-45.
9. Zamfirescu, N., 1977, *Bazele biologice ale producției vegetale*. Ceres Printing House, Bucharest, p.337.
10. Zăhan, P., R., Zăhan, 1989, *Cercetări privind influența plantei premergătoare și a fertilizării asupra dinamicii de acumulare a masei vegetale la grâul cultivat pe soluri podzolice cu exces temporar de umiditate din Câmpia de Vest a țării (I)*. Probleme de agrofitotehnie teoretică și aplicată nr. 1, vol. XI, pp.97-102.
11. Zăhan, P., R., Zăhan, 1989, *Cercetări privind acumularea biomasei vegetale radiculare și calitatea recoltei obținute, sub influența plantei premergătoare și a fertilizării la grâul cultivat pe soluri podzolice cu exces temporar de umiditate din Câmpia de Vest a țării (II)*. Probleme de agrofitotehnie teoretică și aplicată, nr. 1, vol. XI, pp. 237-240.