

## THE INDUCEMENT AT THE ROOTEDNESS PROCESS OF DIERVILA FLORIDA CUTTING USING RADISTIM TYPE BIOACTIVE SUBSTANCE

Vlad Mariana, Vlad Ioan, Vlad Ioana

University of Oradea, Environmental Protection Faculty, 26 General Magheru St.,  
4100848, Oradea, Romania: [ioanvlad2006@yahoo.com](mailto:ioanvlad2006@yahoo.com)

### Abstract

*Diervila florida* is a shrub cultivated as decorative plant for its small, lasting and shining leaves, oval-elongated (5-7 cm/2,5-4,5 cm), with short tail (0,5-1,5 cm) and for the various colored sepals (pink, red, violet) gathered around the yellow flowers with no decorative value. In present it is known as a decorative plant cultivated in field. In areas with less favorable climate conditions where the minimal temperature goes down during winter below the resistance limit, they are cultivate in pots placed outdoor during the summer and indoor during the cold season. In our country *Diervila florida* is no very spread because of the shortage of cutting caused by the low rate of multiplication. In order to increase the efficiency of multiplication on vegetative way, between 2009-2011, in the green houses from Oradea we have watched over the *Diervila florida* cuttings rootedness process using stimulating substances of Radistim type.

**Key words:** *Diervila florida*, rooting substrate, variants, cuttings

### MATERIAL AND METHOD

There where inghathered cuttings semi-wooden 8-10 cm long. The experiment was organized in two variants: V<sub>1</sub> - untreated standard and V<sub>2</sub> - treatment with radistim 2, using 1000 cutting per variant in four different times.

Cutting planting for striking roots has been made in perlite with 1-1,5 mm particles, placed on the parapet with a thickness of substratum of 12-14 cm. The treatment was made before planting. First there was a renewed the humidity status. Then the cuttings were inserted in the powder stimulating substance (radistim 2) with 1-2 cm of their root.

The cutting were planted for striking roots in the first decade of Mai. The distance between cuttings was 6x6 cm and the depth was 2-3 cm. The soil was well ramed in order to remove the air from the rootedness zone.

During the rootedness period the temperature oscillated between 18-27 celsius degrees in air and 20-21 celsius degrees in substratum. The substratum's humidity was 65-75% of total capacity of retaining and the relative humidity was 75-85%.

The light was directed by covering the cuttings with paper and windows of the green house were whitewashed once the growing process started. For the variants differentiation there were made observations and

determinations concerning the length of rootedness period, the proportion of rooted cuttings and the dimensions of new formed roots.

## RESULTS AND DISCUSSIONS

First roots appeared at closed intervals of time for the two variants with a slight advantage for the cuttings treated with Radistim 2.

The period of complete rootedness process lasted 57 days (04.05-30.06).

After the striking root process the cuttings were dislocated from the rootedness substratum and they were passed in clay flower pots which have the diameter of 6-8 cm. In these pots the substratum is formed of: two parts peat, one part earth of leaves, one part compost and one part sand.

The number of rooted cuttings from the total cutting planted for rootedness, for each variant registered growing values from 604 cuttings for V<sub>1</sub> (control, standard variant) to 742 cuttings for V<sub>2</sub> when the cuttings were treated with Radistim 2 (Table 1).

*Table 1*

The striking roots proportion of *Diervila florida* cuttings at Oradea's green houses (average values 2009-2011)

Variants	Number of rooted cutting		±D	Signification of the difference
	Absolute (pcs.)	Relative (%)		
V1 - untreated standard (control variant)	604	100	-	-
V2 - treatment with Radistim	742	122.80	138	xxx

LSD 5% - 6,34

LSD 1% - 8,27

LSD 0.1% - 11,62

In relative terms treatment with Radistim 2 increased the rate of cuttings striking roots with 22% comparatively with the untreated variant. From the statistic point of view this difference is considered as very meaningful.

The treatment with radistim 2 stimulates also the quality of rooted cutting through the number and the dimension of the roots.

From table no.2 arises that the average number of roots per cutting is growing from 8,2 pcs. at V<sub>1</sub>- untreated, to 11,7 pcs. per cutting at V<sub>2</sub>-treated with Radistim 2.

In relative terms the treatment with Radistim 2 increased the number of roots per cutting with 42% comparatively with the untreated variant. From the static point of view this difference is considered as very meaningful.

Table 2

Average number of roots per cutting (average values 2009-2011)

Variants	Average number of roots		±D	Semnification of the diference
	Absolute (pcs.)	Relative (%)		
V1 - untreated standard (control variant)	8,2	100	-	-
V2 - treatment with Radistim	11,7	142.60	3,5	xxx

LSD 5% - 3,45                      LSD 1% - 5,11                      LSD 0.1% - 7,59

The increasing capacity of striking roots arises also from the number and the thickness of the newly formed plants' roots.

From the table no.3 we can see that the lenght and the thickness of *Diervila florida* cutting vary between large limits with favor for those treated with Radistim.

Table 3

The lenght and the thickness of *Diervila florida* rooted cuttings (average values 2009-2011)

Variants	The lenght of roots-extreme limits (cm)	Grouping the roots in accordance with its thickness		Total
		Pes.<1mm	Prs.>1mm	
V1 - untreated standard (control variant)	0,4-9,5	4,7	3,5	8,2
V2 - treatment with Radistim	0,5-12,7	6,9	4,8	11,7

For the control variant the newly formed roots registered variable lenght between 0,4 and 9,5 cm. For the cutting treated with radistim 2 the value were higher, between 0,5 and 12,7 cm.

Grouping the newly formed roots in accordance with its thickness, for the roots with diameter smaller that 1 mm there were registered values in growth from 4,7 pcs. for V<sub>1</sub> to 6.9 pcs. for V<sub>2</sub>. For the roots with diameter bigger that 1 mm there registered values in growth from 3,5 pcs. for V<sub>1</sub> to 4,8 pcs. for V<sub>2</sub>.

## CONCLUSIONS

\* *Diervila florida* as decorative species, with useful economic implications, can be multiplied through vegetative way by cuttings.

\* The multiplication rate of *Diervila florida* through cuttings can be stimulated by using bioactive substances of Radistim type.

\* Stimulating the rootedness process of semiwooden cutting of *Diervila florida* with bioactive substances of Radistim type guarantee a highly vegetative potential for newly formed plants.

\* The stimulate substance Radistim increase the striking roots rate. So the treated cuttings strike roots in proportion of 74,2% comparatively to 60,4% for those untreated.

## REFERENCES

1. Andekerk, Th. G. L. (1990). Salt tolerance of ten woody ornamentales. Abstr. Of the XXXIII<sup>th</sup> International Horticulture Congress, Firenze.
2. Alborede, M.(1992). La mise en bac d'un arbre adulte. Revue Horticole, avril.
3. Boutherin, D. and Bron G. (1999). Multiplication des plantes horticols, Ed. Tehnique et Documentation Lavoisier, Paris.
4. Bush-Brown. (1995). Garden Book, Charrles Scribners's Sons, USA.
5. Contet, A. (1999). Pepiniere d'ornament et fruitiere, Ed. Bailliere et Filis, Paris.
6. Cuisance, P.(1992). Les arbustes d'ornament, Ed. Floraisse Larouse, Paris.
7. Enescu, V. and Ioniță L. (1994). Paladă Nicolau M.- Înmulțirea vegetativă a arborilor forestieri, Ed. Ceres, București.
8. Foucard, J. (2004). Filiere Pepiniere, TEC-DOC Lavoisier, Paris.
9. Gorastarzu, B. (2002). Bacetrisation des substrates et mycorhization centrale, L'horticulture Francaise, mars.
10. Harris, W. (2006). Arboriculture syllabus. Department of Enviromental Horticulture, University of California Davis, USA.
11. Hay, R. and Synage P. (2007). 2000 fleurs, plantes et arbustes, Ed Oyez, Leuven, Belgique.
12. Herwing, R. (2008). L'enciclopedie pratique des fleurs, plantes et arbres de jardin, Ed. Culture, Art, Loisir, Paris.
13. Hessayeon, D. (2003). The Tree and Shrub Expert, Publication Britanica House, Great Britain.
14. Iliescu, A. F. (1998). Arboricultură ornamentală, Ed. Ceres, București.
15. Jhenesen, H. (2004). Le grand livre international des arbres, Ed. F. Nathan, Paris.
16. Krussman, G, (1998). Die Laubgehölze. Verlag Paul Parey, Berlin.
17. Krussman, G, (2005). Die Laubgehölze. Verlag Paul Parey, Berlin
18. Nitschke, Gunter (1993). Le jardin japonais, Taschen, Koln.
19. Mailliet, L. (2003). Arboriculture urbaine, Institute Pour le development forestier, Paris.
20. Thome, S. (2002). Pepiniere d'arboriculture ornamentale 151, Gremblause.
21. Vlad, I. (2004). Floriculture, Imprimeria de Vest Publishing house, Oradea.