

## THE INFLUENCE OF CROP ROTATION AND NUTRITION REGIME ON THE QUALITY INDICATORS OF SEEDS IN WINTER WHEAT CULTIVATED ON THE PRELUVOSOILS

Bandici Gheorghe\*, Ardelean Ileana\*, Bucurean Elena\*

\*University of Oradea, Faculty of Environmental Protection, 26 General Magheru St., 410048  
Oradea, Romania, e-mail: [gbandici@yahoo.com](mailto:gbandici@yahoo.com)

### **Abstract**

*The quality of production is related to a series of physical and chemical characteristics of plants which gives a positive mark to the applied agrotechnical methods for the correlation of the latter to the production obtained on the surface unit.*

**Key words:** crop rotation, nutrition regime, nitrogen, phosphorus, potassium., raw protein, seeds, winter wheat

### **INTRODUCTION**

The production quality is related to a series of physical and chemical characteristics of the plants which gives a positive mark to the agrotechnical applied measures for the correlation of this with the production obtained for the surface unit (Munteanu, et al., 2011, Domuța, 2012).

The research performed in this field made clear the fact that quality is conditioned by the species and the cultivated hybrid, the climatic conditions of the cultivating year and also by the technology applied to the agricultural plants (Dincă, 1982, Bilteanu, 1993). To justify some of these aspects with consequences regarding the quality of the final production, we make some references to the specialised scientific literature, i.e. Hera Cr. and her team (1986) underline the importance of nitrogen for the increase of the protein content, wet and dry gluten and for the improvement of the quality indicators of gluten. The authors also mention the importance of the ameliorative plant (the pea) for the quality indicators of the wheat. Boldea Eleonora and her team (1986) also mention the importance of the new species of wheat for the quality of raw protein and gluten.

The main component of the chemical composition of the seeds is represented by the glucides (62-75 %) of the fresh wheat grain mass, the proteins 10-16 %, lipids 1.8-2.6 %, cellulose 2-3.5 % and mineral substances 1.5-2.3 % (Hera, 1986, Soltner, 1990, Salisbury, 1995). A series of analyses of the N, P, K and raw protein content in the wheat grains has been made in order to specify the quality of the final product (Bandici, 1997; 2001, Ardelean 2006, 2013).

Some analyses have been made to establish the quality of the final product regarding the content of N, P, K in wheat seeds and raw protein (Bandici, and all., 2003, Domuța and al., 2007, 2008).

## MATERIAL AND METHODS

The experiment was made at Agricultural Research and Development Station (A.R.D.S) Oradea, in the period 2016-2018, on the preluvosoils. For “Delia” winter wheat grains a series of chemical test were made regarding the content of nitrogen, phosphorus, potassium and raw protein according to the precursory and the nutrition system. The nitrogen was determined using the Kjeldahl method, the phosphorus was determined by colorimetry with ammonium molybdate and tin chloride reduction. The potassium was determined through flame photometry and the raw protein was determined through calculation ( $N_t \times 5.7 \%$ ).

## RESULTS AND DISCUSSION

Analysing the data in table 1, regarding the influence of forerunner plant and fertilization level on the total N content in the wheat seeds, we can see that both the forerunner plant and fertilization level influenced the content of this element in seeds. There fore, comparing the wheat monoculture with wheat cultivation that was preceded by corn = maize or pea (3 and 4 years crop rotation) the latter induces an increased production of 22.4-53.8 %.

As an ameliorative plant, pea determined the increase of nitrogen content in the crop as a consequence of its symbiotic particularities. Compared to the unfertilized type, with a value of 1.37 g/100 g.d.w. (grains of dry substance = wheat), mineral and organo-mineral fertilization determine important increase of nitrogen, i.e. 38.7 % and 62 %.

Table 1

The influence of crop rotation and nutrition regime of the final content of *nitrogen* of the seeds in wheat cultivated on preluvosoils, Oradea, 2016-2018

Observed factor	Total g/100 g.d.w.	Nitrogen %		Difference +/-
a.	b. Crop rotation			
Wheat – Monoculture (C <sub>1</sub> )	1.43	100		-
Maize (W-M)	1.75	122.4		+0.32
Pea (P-W-M)	2.20	153.8		+0.77
Pea (P-W-M-M)	1.95	136.4		+0.52
c.	d. Nutrition regime			
N <sub>0</sub> P <sub>0</sub> (C <sub>1</sub> )	1.37	100		-
N <sub>120</sub> P <sub>80</sub>	1.90	138.7		+0.53
N <sub>120</sub> P <sub>80</sub> +10 t/ha manure	2.27	162.0		+0.85

In point of the factors interaction: crop rotation x nutrition regime (table 2), we note that no matter the crop rotation used, mineral or organo-mineral fertilization increase by 12.1-86.7 %. The lowest values of total nitrogen content can be found in the wheat monoculture (1.24-1.65 g/100 g.d.w.) compared to short wheat – maize rotation (1.27-2.07 g/100 g.d.w.) or to 3 and 4 year wheat – pea crop rotation – (1.70-2.78 g/100 g.d.w. and 1.28-2.39 g/100 g.d.w.).

Table 2

Influence of the factors interaction: crop rotation x nutrition regime on the final content of nitrogen of the seeds in wheat cultivated on preluvosoils, Oradea 2016-2018

Nutrition regime	Total g/100 g.d.w.	Nitrogen %	Difference +/-
a. Wheat – Monoculture (M <sub>t</sub> )			
N <sub>0</sub> P <sub>0</sub> (C <sub>t</sub> )	1.24	100	-
N <sub>120</sub> P <sub>80</sub>	1.39	112.1	+0.15
N <sub>120</sub> P <sub>80</sub> +10 t/ha manure	1.65	133.1	
b. Maize (W-M)+0,41			
N <sub>0</sub> P <sub>0</sub> (M <sub>t</sub> )	1,27	100	-
N <sub>120</sub> P <sub>80</sub>	1.90	149.6	+0.63
N <sub>120</sub> P <sub>80</sub> +10 t/ha manure	2.07	163.0	+0.80
c. Pea (P-W-M)			
N <sub>0</sub> P <sub>0</sub> (M <sub>t</sub> )	1.70	100	-
N <sub>120</sub> P <sub>80</sub>	2.13	125.3	+0.43
N <sub>120</sub> P <sub>80</sub> +10 t/ha manure	2.78	163.5	+1.08
d. Pea (P-W-M-M)			
N <sub>0</sub> P <sub>0</sub> (M <sub>t</sub> )	1.28	100	-
N <sub>120</sub> P <sub>80</sub>	2.18	170.3	+0.90
N <sub>120</sub> P <sub>80</sub> +10 t/ha manure	2.39	186.7	+1.11

Concerning the *total raw protein* content (Nt x 5.7), in the table 3 and 4 we note the direct link between the nitrogen content and raw protein.

In this case, the crop rotation and the nutrition regime in the process induce important raw protein increase, which, in case of 3 year wheat-pea crop rotation may rise up to 12.58 g/100 g.d.w., compared to monoculture of 8.15 g/100 g.d.w. The highest values of raw protein increase were established in the organo-mineral fertilization process of 12.58g/100 g.d.w., compared to the witness (N<sub>0</sub>,P<sub>0</sub>) 7.92 g/100 g.d.w. In the case of raw protein, no matter what the precursory was, the organo-mineral fertilization determined the highest values of raw protein content which varied between 9.43 g/100 g.d.w., in wheat monoculture and 15.84 g/100 g.d.w., in pea (3 year crop rotation).

Table 3

The influence of crop rotation and nutrition regime on the final content of *raw protein* of the seeds in wheat cultivated on preluvosoils, Oradea 2016– 2018

Observed factor	Raw protein g/100 g.d.w.	Raw protein %	Difference +/-
a. Crop rotation			
Wheat – Monoculture (M <sub>t</sub> )	8.15	100	-
Maize (W-M)	9.96	118.5	+1.81
Pea (P-W-M)	12.58	154.3	+4.43
Pea (P-W-M-M)	11.23	137.8	+3.08
b. Nutrition regime			
N <sub>0</sub> P <sub>0</sub> (M <sub>t</sub> )	7.92	100	-
N <sub>120</sub> P <sub>80</sub>	10.84	136.9	+2.92
N <sub>120</sub> P <sub>80</sub> +10 t/ha manure	12.68	160.1	+4.76

Table 4

Influence of the factors interaction: crop rotation x nutrition regime on the final content of *raw protein* of the seeds in wheat cultivated on luvosoils, Oradea 2016-2018

Observed factor	Raw protein g/100 g.d.w.	Raw protein %	Difference +/-
a. Wheat – Monoculture (M <sub>t</sub> )			
N <sub>0</sub> P <sub>0</sub> (M <sub>t</sub> )	7.07	100	-
N <sub>120</sub> P <sub>80</sub>	7.95	112.4	+0.88
N <sub>120</sub> P <sub>80</sub> +10 t/ha manure	9.43	133.3	+2.36
b. Maize (W-M)			
N <sub>0</sub> P <sub>0</sub> (M <sub>t</sub> )	7.26	100	-
N <sub>120</sub> P <sub>80</sub>	10.83	149.2	+3.57
N <sub>120</sub> P <sub>80</sub> +10 t/ha manure	11.79	162.4	+4.53
c. Pea (P-W-M)			
N <sub>0</sub> P <sub>0</sub> (M <sub>t</sub> )	9.72	100	-
N <sub>120</sub> P <sub>80</sub>	12.17	125.2	+2.45
N <sub>120</sub> P <sub>80</sub> +10 t/ha manure	15.84	163.1	+6.12
d. Pea (P-W-M-M)			
N <sub>0</sub> P <sub>0</sub> (M <sub>t</sub> )	7.62	100	-
N <sub>120</sub> P <sub>80</sub>	12.43	163.1	+4.81
N <sub>120</sub> P <sub>80</sub> +10 t/ha manure	13.65	179.1	+6.03

Regarding the total content of *phosphorus* in the wheat seeds, in *table 5* we note that neither crop rotation, nutrition regime, nor their interaction led to significant differences, regardless of the quality of the forerunner plant or organo-mineral fertilization, except the pea (3 year crop rotation) when the mineral or organo-mineral fertilisation determined more than 10 % increase of the total content of phosphorus.

*Table 5*

Influence of the crop rotation and nutrition regime on the final content of *phosphorus* of the seeds in wheat cultivated on preluvosoils, Oradea 2016-2018

Observed factor	Total phosphorus g/100 g.d.w.	Phosphorus %	Difference +/-
a. Crop rotation			
Wheat – Monoculture (M <sub>i</sub> )	0.36	100	-
Maize (W-M)	0.36	100	-
Pea (P-W-M)	0.40	111.0	+0.04
Pea (P-W-M-M)	0.36	100	-
b. Nutrition regime			
NoP <sub>0</sub> (M <sub>i</sub> )	0.36	100	-
N <sub>120</sub> P <sub>80</sub>	0.37	102.8	+0.01
N <sub>120</sub> P <sub>80</sub> +10 t/ha manure	0.38	105.5	+0.02

Regarding the total content of *potassium* in the wheat seeds, in *Table 6* under the individual influence of both the observed factors, we could notice significant difference.

*Table 6*

Influence of the forerunner plant and fertilization level on the final content of *potassium* of the seeds in wheat cultivated on preluvosoils, Oradea 2016-2018

Observed factor	Total g/100 g.d.w.	Potassium %	Difference +/-
a. Crop rotation			
Wheat – Monoculture (M <sub>i</sub> )	0.64	100	-
Maize (W-M)	0.67	104,7	+0.03
Maize (P-W-M)	0.64	100	-
Pea (P-W-M-M)	0.63	98,0	+0.01
b. Nutrition regime			
NoP <sub>0</sub> (M <sub>i</sub> )	0.67	100	-
N <sub>120</sub> P <sub>80</sub>	0.63	94.0	-0.04
N <sub>120</sub> P <sub>80</sub> +10 t/ha manure	0.63	94.0	-0.04

## CONCLUSIONS

A more intense accumulation of the biomass which determines an intensification of the photosynthesis positively influences the chemical composition of the final product – the grains.

The total content of nitrogen in the winter wheat grains was influenced by the crop rotation and the nutrition system.

The raw protein content follows the natural way similarly to nitrogen total content being influenced mainly by the crop rotation and the fertilization level.

There weren't observed any essential changes of the total phosphorus and potassium content under the influence of the crop rotation and the fertilization level.

## REFERENCES

1. Ardelean Ileana, 2006, Contribution in the known and modification of the crop rotation influence on quantity and quality of the winter wheat yield cropped on the acid soils from North-Western Romania. Thesis degree, USAMV Cluj-Napoca, p.220.
2. Ardelean Ileana, 2013, Agrotehnica. Editura Universității din Oradea. p.417
3. Bandici G. E., 1997, Contributii la stabilirea influenței premergătoarei și a fertilizării asupra dinamicii acumulării biomasei, la grâul de toamna, cultivat pe soluri cu exces temporar de umiditate, în centrul Câmpiei de Vest a României. Doctoral thesis. University of Agriculture Sciences and Veterinary Medicine Cluj-Napoca, Romania in Romanian, p.250.
4. Bandici G. E., C. Domuta, Ileana Ardelean, 2003, The influence of the forerunner plant, fertilisation level and climatic conditions on the total wet and dry gluten content of winter wheat seeds cultivated on brown luvisc soils in the Western Plain of Romania. Lucrari științifice USAMVB., Seria B, vol. XLV, Bucuresti, p.281-284, p.330.
5. Bandici, G.E., Guș, P., 2001, Dinamica acumulării de biomasă la grâul de toamnă. University of Oradea Press, 107 p.
6. Bîlteanu, G., 1993.: Fitotehnie, Ceres Printing House. Bucharest, pp. 457.
7. Boldea, Elena, 1986, Însușirile de panificație a unor soiuri de grâu raionate și linii de perspectivă, Probleme Agricole, nr.7, p.27-32, p.50.
8. Dincă, D., 1982: Asolamentele agriculturii moderne. Ceres Printing House. Bucharest. 257 pp
9. Domuta C., 2012, Agrotehnică. Editura Universității din Oradea. p.506
10. Domuta C., Bandici Gh., Ciobanu Gh., N. Csep, Ciobanu Cornelia, Samuel Alina, Bucurean Elena, Sandor Maria, Borza Ioana, Bunta Gh., Ileana Ardelean, Cr. Domuta, 2007, "Asolamentele în Câmpia Crișurilor". Editura Universității din Oradea, ISBN 978-973-759-350-4, pag. 254.
11. Domuta C. , Bandici Gh., Ciobanu Gh. Ciobanu Cornelia, Samuel Alina, N. Csep, Bucurean Elena, Borza Ioana, Sandor Maria, Bunta Gh., Ileana Ardelean, Cr. Domuța., 2008, "Asolamentele in sistemele de agricultura", Editura Universității din Oradea. ISBN , pag. 297..
12. Hera, C., 1986, Influența fertilizării asupra unor indici calitativi ai recoltelor de grâu, Probleme de agrofitehnie teoretică și aplicată, nr.2, vol.VIII, p.71-76, p.100.
13. Hera, C., 1986, Influența unor factori tehnologici asupra calității grâului, Cereale și plante tehnice, nr.7, p.47-52, p.88.
14. Muntean L. S. S., G. Cernea, G. Morar, M. Duda, D. Vârban, S. Muntean, 2008, Fitotehnie. Academic Press Printing House, Cluj-Napoca, p.83-135, p.255.
15. Salisbury F.B., C.W. Ross, 1995 - Fisiologia vegetale. Seconda edizione italiana condota sulla quarta edizione americana. Editura Zanichelli.
16. Soltner D., 1990, „Phytotechnie speciale”, Collection sciences et Techniques Agricoles, Angers.