

PHYTOCOENOLOGICAL STUDY OF ROCKY VEGETATION FROM CODRU-MOMA MOUNTAINS BASED ON ECOLOGICAL INDICES

Pășcuț Călin Gheorghe*

* University of Oradea, Faculty of Environmental Protection, 26 Gen. Magheru St., 410048,
Oradea, Romania, e-mail: pascutcalin@yahoo.com

Abstract

In this paper the ecological factors for the rocky associations from Codru-Moma Mountains are studied (humidity, temperature, chemical reaction of soil). The study was done during 2008-2018, a number of 5 rocky associations being identified: *Asplenietum rutae-murariae-trichomanis* R. Tüxen 1937 (25 relevées), *Asplenio quadrivalenti-Poëtum nemoralis* Soó ex Gergely et al. 1966 (17 relevées), *Asplenietum septentrionali-adianti-nigri* Oberdorfer 1938 (9 relevées), *Asplenio trichomani-Poëtum nemoralis* Boșcaiu 1971 (25 relevées), *Ctenidio-Polypodietum* Jurko et Peciar 1963 (25 relevées).

In Codru-Moma Mountains the phytocoenoses of these associations have been identified at altitudes of 330-950 m, on siliceous or calcareous rocks, with large inclination (30-90°) and shaded exposition (N, E, V, NV, NE), along some valleys and streams on the edge of hornbeam and beech forests. Vegetation covers 25-100%, and due to the fact that on these cliffs humidity is high all year round, a series of bryophytes appear (1-40%).

Key words: ecological factors, humidity, temperature, chemical reaction of soil, relevées

INTRODUCTION

The rocky phytocoenoses represent the pioneer vegetation that develops in the cracks of calcareous and siliceous rocks from Codru-Moma Mountains. Due to the nature of the substrate and the restrictive climatic conditions, the floristic composition of these associations is characterized by an abundance of species of *Quercus-Fagetum*, *Festuco-Brometum*, *Rhamno-Prunetum* classes.

The main plant rocky associations identified in the Codru-Moma Mountains and analyzed in this paper from the ecological point of view are:

Asplenietum trichomanis Class (Br.-Bl. in Meier et Br.-Bl. 1934)
Oberdorfer 1977

Tortulo-Cymbalarietalia order Segal 1969

Cymbalario-Asplenion alliance Segal 1969 em. Mucina 1993

1. *Asplenietum rutae-murariae-trichomanis* association R. Tüxen 1937

Cystopteridion alliance Richard 1972

2. *Asplenio quadrivalenti-Poëtum nemoralis* association Soó ex Gergely et al. 1966

Androsacetalia vandelli order Br.-Bl. in Maier et Br.-Bl. 1934

Asplenion septentrionalis alliance Oberdorfer 1938

3. *Asplenietum septentrionali-adianti-nigri* association Oberdorfer 1938

4. *Asplenio trichomani-Poëtum nemoralis* association Boşcaiu 1971

Hypno-Polypodium alliance Mucina 1993

5. *Ctenidio-Polypodietum* association Jurko et Peciar 1963

Older studies related to the rocky vegetation from Codru-Moma Mountains were made by Paucă (1941). More recent publications relating to the rocky vegetation from Codru-Moma Mountains were made by Pășcuț (2010, 2013), Burescu and Pășcuț (2010).

In our country and in Europe, these rocky associations were studied by Pascal and Mititelu (1971), Pop et al., (2002), Rațiu et al., (1984), Ștefan et al., (1997), Sârbu et al., (1997), Chifu et al., (2006), Chifu et al., (2014), Oprea and Sîrbu (2009), Rațiu and Gergely (1976), Schneider-Binder (1968, 1969, 1972), Mihăilescu (2001), Karácsonyi, (2011), Groza, (2008), Niculescu (2006), Stancu (2005), Răduțoiu (2006), Drăgulescu (1985, 1988), Diaconescu (1973), Alexiu (1998), Peia, (1978), Kolbek et al., (2015), Pott (1995), Borhidi (2003), Pignatti and Pignatti (2014), Świerkosz (2004).

MATERIAL AND METHOD

For the study of the identified associations we used ecological spectra that take into account the presence of the species in the association, as well as humidity, temperature and chemical reaction of soil for each species (Sanda et al., 1983). The phytocoenological description of the associations is followed by comparative analysis in order to outline the requirements of each association on ecological factors.

In the table constancy (K) is mentioned for each identified species. The constancy (K) outlines the general presence of a species in all the relevés analysed, belonging to the same type of phytocoenose (Cristea et al., 2004).

In the reports of the relevés information regarding altitude, herbaceous layer coverage, bryophytes layer coverage, exposition and slope was introduced.

Information concerning life forms, floristic elements and ecological indices is given based on Sanda et al., (1983), Cristea et al., (2004).

Nomenclature adopted for the identified species is in accordance with the work developed by the Ciocârlan (2009) și Sârbu et al., (2013).

RESULTS AND DISCUSSION

The rocky associations identified in the Codru-Moma Mountains are presented as follows:

1. *Asplenietum rutae-murariae-trichomanis* association R. Tüxen 1937

The phytocoenose of *Asplenietum rutae-murariae-trichomanis* association (figure 1), develops in the form of clumps among the cracks and on shelves of heavily skewed limestone rocks (30-90°), mainly with north, northwest or northeast exposition. It is widespread on the rocks along the intermountain valleys and creeks at altitudes of 400-680 m.

Thanks to the nordic exhibition of the rocks and the inclusion of these phytocoenoses in beech forests, a shaded microclimate with moderate humidity almost all year round has been created, as well as lower temperature in summer. This has allowed the installation of bryophytes of the genus *Ctenidium* and *Syntrichia* within these phytocoenoses.

Considering life forms, the hemycryptophytes (59.37%) dominate, followed by terophytes (17.19%) and chamaephytes (10.94%).

The most numerous floristic elements are eurasian (35.94%), followed by european (20.31%), central-european (15.63%) and mediterranean (7.8%) ones.

2. *Asplenio quadrivalenti-Poëtum nemoralis* association Soó ex Gergely et al. 1966

The association phytocoenoses vegetate on shaded limestone rocks, with high slope (40-90°), located at altitudes of 410-800 m, set on shaded slopes (N, NV, NE). The physiognomy of the association is given by the two edifying species *Asplenium trichomanes* ssp. *quadrivalens* and *Poa nemoralis*, in terms of codominance.

The life forms are dominated by hemycryptophytes (52.38%), followed by phanerophytes (23.8%), geophytes and terophytes (11.91%) having an equal share.

The floristic elements show the predominance in the association of the eurasian species (38.09%), followed by european (23.81%), central-european (16.67%), cosmopolitan (9.53%) and south european (4.76%) ones.

In this association the bryophytes layer consisting of *Ctenidium molluscum* and *Marchantia polymorpha* plays an important role in maintaining moisture.

3. *Asplenietum septentrionali-adianti-nigri* association Oberdorfer 1938

In Codru-Moma Mountains the phytocoenoses of this association (figure 2), have been identified at altitudes of 350-590 m, on siliceous rocks with great slope (40-80°), along some valleys and streams on the edge of hornbeam and beech forests. They develop on superficial soil, poor in organic substances and with a low acid neutrophile PH.

The life forms outline the presence of a large number of hemycryptophyte species (75%), followed by chamaephytes (10.71%) and nanophanerophytes (7.14%).

Floristic elements with the largest share in the association are the eurasians (42.86%), followed by european species (14.29%), central-european species (10.71%), cosmopolitan (10.71%) and circumpolar (10.71%) ones.

On these rocks there is high humidity all year round, demonstrated by the presence of *Ctenidium molluscum* species.



Fig. 1. *Asplenietum rutae-murariae-trichomanis* (Râposu Brook, Bihor County)



Fig. 2. *Asplenietum septentrionali-adianti-nigri*, (Briheni Valley, Bihor County)

4. *Asplenio trichomani-Poëtum nemoralis* association Boşcaiu 1971

It is an association found on siliceous rocks, with great slope (30-90°), at altitudes between 350-945 m. The phytocoenoses of this association (figure 3), prefer the rocks in shady valleys, located on slopes with northern and north-western exposition.

Considering life forms hemycryptophytes (52.38%) have the highest share, followed by the phanerophytes (25.39%), terophytes (12.7%) and geophytes (9.52%).

Among the floristic elements in the phytocoenoses analyzed the eurasian species (34.92%) dominate, supplemented by european (19.05%), central-european (17.46%) and south european (9.53%) ones.

Bryophytes layer is well represented, consisting of *Ctenidium molluscum*, *Hypnum cupressiforme*, *Marchantia polymorpha*, *Polytrichum commune*.

An important feature is a large fluctuation of coincidental species generated by the variability of vegetation conditions.

5. *Ctenidio-Polypodietum* association Jurko et Peciar 1963

The *Ctenidio-Polypodietum* phytocoenoses (figure 4), develop in the cracks and on shaded rocks, on lower hills and up to the top of Codru-Moma Mountains, at altitudes of 330-1000 m, on terrains with a shady exposition (N, NV, NE) and high slope (30-90°). Humidity is the determining factor, the association being mainly found on the lower third of steep and shady slopes, near the mountain valleys and creeks.

In the case of life forms one can observe the presence of the large number of hemycryptophyte species (57.81%), followed by phanerophytes (17.19%), geophytes (12.5%) and terophytes (12.5%).

Considering floristic elements, you can see the share of eurasian species (26.56%), european ones (23.44%), followed by central-european (15.62%), cosmopolitan (12.5%) and south european (9.37%) species.



Fig. 3. *Asplenio trichomanis*-*Poëtum nemoralis* (Urviş Valley, Arad County)



Fig. 4. *Ctenidio-Polypodietum* (Finiş Valley, Bihor County)

The following table presents in detail the floristic composition of the associations identified, as well as the constancy (K) for each species.

Indices of species constancy for the analyzed associations					Table 1
The number of the association	1	2	3	4	5
Altitude (m.s.m.)	400-680	410-800	350-590	350-945	330-950
The coverage of vegetation layer (%)	25-80	30-85	40-80	40-80	50-100
The coverage of bryophytes layer (%)	1-25	5-40	5-20	1-25	1
Exposition	N, NE, NV	N, NE, NV, E	N, V, NV	N, V, E, NV, NE	N, V, NE, NV
Slope (°)	30-90	40-90	80-90	30-90	30-90
	0	1	2	3	4
<i>Asplenietea trichomanis</i>					
<i>Asplenium trichomanes</i> ssp. <i>quadrivalens</i>	V	V	.	.	II
<i>Asplenium trichomanes</i> ssp. <i>trichomanes</i>	.	.	V	V	III
<i>Asplenium ruta-muraria</i>	V	I	.	.	I

0	1	2	3	4	5
<i>Asplenium septentrionale</i>	·	·	V	I	I
<i>Asplenium adiantum-nigrum</i>	·	·	V	I	I
<i>Poa nemoralis</i>	II	V	IV	V	III
<i>Sedum maximum</i>	II	III	V	IV	IV
<i>Polypodium vulgare</i>	II	II	II	IV	V
<i>Valeriana tripteris</i>	II	II	·	I	I
<i>Cystopteris fragilis</i>	I	II	·	I	I
<i>Sile nutans</i> ssp. <i>dubia</i>	·	·	II	·	·
<i>Cardaminopsis arenosa</i>	I	I	·	I	I
<i>Asplenium adulterinum</i>	·	I	I	·	I
<i>Sedum hispanicum</i>	I	·	·	·	·
<i>Calamagrostis arundinacea</i>	·	·	·	·	I
<i>Ceterach officinarum</i>	·	·	·	·	I
Festuco-Brometea	·	·	·	·	·
<i>Teucrium chamaedrys</i>	II	·	III	·	·
<i>Sedum acre</i>	I	·	III	·	·
<i>Thymus glabrescens</i>	I	·	III	·	·
<i>Euphorbia cyparissias</i>	I	·	II	·	·
<i>Hypericum perforatum</i>	I	·	II	·	·
<i>Potentilla argentea</i>	I	·	II	·	·
<i>Achillea collina</i>	I	·	·	·	·
<i>Agrimonia eupatoria</i>	I	·	·	·	·
<i>Asperula cynanchica</i>	I	·	·	·	·
<i>Brachypodium pinnatum</i>	I	·	·	·	·
<i>Centaurea biebersteinii</i>	I	·	·	·	·
<i>Dianthus carthusianorum</i>	I	·	·	·	·
<i>Festuca valesiaca</i>	I	·	·	·	·
<i>Fragaria viridis</i>	I	·	·	·	·
<i>Geranium pusillum</i>	I	·	·	·	·
<i>Poa compressa</i>	I	·	·	·	·
<i>Potentilla arenaria</i>	I	·	·	·	·
<i>Sanguisorba minor</i>	I	·	·	·	·
<i>Sedum sexangulare</i>	I	·	·	·	·
<i>Stachys germanica</i>	I	·	·	·	·
<i>Teucrium montanum</i>	I	·	·	·	·
<i>Thymus comosus</i>	I	·	·	·	·
<i>Thymus pannonicus</i> ssp. <i>pannonicus</i>	I	·	·	·	·
<i>Trifolium arvense</i>	I	·	·	·	·
<i>Viola arvensis</i>	I	·	·	·	·
<i>Vincetoxicum hirundinaria</i>	I	·	·	·	·
Quercu-Fagetea					
<i>Geranium robertianum</i>	II	III	II	III	IV
<i>Lamium galeobdolon</i>	II	III	·	III	IV
<i>Asplenium scolopendrium</i>	III	IV	·	II	II
<i>Rubus hirtus</i>	·	II	·	IV	III
<i>Mycelis muralis</i>	II	III	II	III	III
<i>Moehringia muscosa</i>	II	III	II	II	II
<i>Dryopteris filix-mas</i>	I	I	II	III	II
<i>Hedera helix</i>	III	II	·	I	II
<i>Cruciata glabra</i>	·	·	III	I	·
<i>Campanula persicifolia</i>	II	I	II	II	II
<i>Asarum europaeum</i>	I	I	·	II	II
<i>Oxalis acetosella</i>	I	·	·	II	II
<i>Galium schultesii</i>	I	II	·	I	I
<i>Polygonatum odoratum</i>	·	I	·	I	II
<i>Athyrium filix-femina</i>	·	·	·	I	II
<i>Galium odoratum</i>	·	·	·	I	II
<i>Luzula luzuloides</i>	·	·	·	I	II
<i>Polystichum aculeatum</i>	·	II	·	·	I

0	1	2	3	4	5
<i>Clinopodium vulgare</i>	.	.	II	.	.
<i>Campanula rapunculoides</i>	I	I	.	I	I
<i>Moehringia trinervia</i>	I	I	.	I	I
<i>Daphne mezereum</i>	I	I	.	I	.
<i>Euonymus latifolius</i>	.	I	.	I	I
<i>Acer pseudoplatanus</i>	.	.	.	I	I
<i>Alliaria petiolata</i>	.	.	.	I	I
<i>Carpinus betulus</i>	.	.	.	I	I
<i>Hepatica nobilis</i>	I	I	.	.	.
<i>Lamium maculatum</i> ssp. <i>maculatum</i>	.	.	.	I	I
<i>Polystichum setiferum</i>	.	.	.	I	I
<i>Stellaria holostea</i>	.	.	.	I	I
<i>Brachypodium sylvaticum</i>	.	.	.	I	.
<i>Carex pilosa</i>	.	.	.	.	I
<i>Dentaria bulbifera</i>	.	.	.	.	I
<i>Fagus sylvatica</i>	.	I	.	.	.
<i>Festuca drymeja</i>	.	.	.	.	I
<i>Glechoma hirsuta</i>	.	.	.	I	.
<i>Pulmonaria officinalis</i>	.	.	.	.	I
<i>Sanicula europaea</i>	.	.	.	I	.
<i>Tilia platyphyllos</i>	.	.	.	.	I
<i>Viola reichenbachiana</i>	.	.	.	I	.
Rhamno-Prunetea					
<i>Sambucus nigra</i>	I	.	.	II	III
<i>Corylus avellana</i>	.	I	.	I	II
<i>Galeopsis speciosa</i>	.	I	.	I	II
<i>Clematis vitalba</i>	.	I	.	I	.
<i>Acer campestre</i>	.	.	.	I	I
<i>Cornus sanguinea</i>	.	.	.	I	.
<i>Crataegus monogyna</i>	.	.	.	I	.
<i>Evonymus europaeus</i>	.	.	.	I	.
<i>Salix caprea</i>	.	.	.	I	.
<i>Ulmus glabra</i>	.	.	.	.	I
<i>Verbascum nigrum</i>	.	.	.	I	.
Variae Syntaxa					
<i>Salvia glutinosa</i>	II	I	II	I	III
<i>Fragaria vesca</i>	.	I	III	I	.
<i>Doronicum austriacum</i>	II	I	.	.	II
<i>Galium album</i>	II	I	.	I	I
<i>Chamaecytisus hirsutus</i> ssp. <i>leucotrichus</i>	.	.	II	I	.
<i>Chelidonium majus</i>	.	.	.	I	II
<i>Genista ovata</i>	.	.	II	.	.
<i>Arabis turrata</i>	I	.	.	I	I
<i>Melampyrum bihariense</i>	.	I	.	I	I
<i>Tamus communis</i>	.	.	.	I	I
<i>Urtica dioica</i>	.	.	.	I	I
<i>Vincetoxicum hirundinaria</i>	.	I	.	.	I
<i>Achillea millefolium</i>	I	.	.	.	.
<i>Cytisus nigricans</i>	.	.	.	I	.
<i>Dianthus spiculifolius</i>	I	.	.	.	.
<i>Festuca pallens</i>	.	.	.	.	I
<i>Galium mollugo</i>	I	.	.	.	.
<i>Gentiana asclepiadea</i>	.	.	.	I	.
<i>Hieracium pilosella</i>	I	.	.	.	.
<i>Hieracium umbellatum</i>	.	.	.	.	I
<i>Medicago lupulina</i>	I	.	.	.	.
<i>Melittis melissophyllum</i>	.	.	.	.	I
<i>Peucedanum longifolium</i>	I	.	.	.	.

0	1	2	3	4	5
<i>Primula elatior</i> ssp.	I	.	.	.	.
<i>leucophylla</i>	I	.	.	.	.
<i>Rosa canina</i>	.	.	.	.	I
<i>Sedum cepaea</i>	.	I	.	.	.
<i>Sambucus nigra</i>	.	.	.	I	.
<i>Solidago virgaurea</i>	.	I	.	.	.
<i>Valeriana officinalis</i>	I	.	.	.	.
<i>Verbascum phlomoides</i>	.	.	.	I	.
<i>Veronica chamaedrys</i>	.	.	.	.	.
Bryophyta					
<i>Ctenidium molluscum</i>	III	IV	IV	II	V
<i>Marchantia polymorpha</i>	.	I	.	I	I
<i>Hypnum cupressiforme</i>	.	.	.	I	.
<i>Polytrichum commune</i>	.	.	.	I	.
<i>Syntrichia ruralis</i> var.	I	.	.	.	.
<i>calcicola</i>					

Place and date of relevés: 1 - *Asplenietum rutae-murariae-trichomanis* association, Brook of Râposu (7 relevés), Morilor Valley (5 relevés), Ormanu Valley (2 relevés), Briheni Valley (2 relevés), Șopotesei Valley, Câmp Moți village (2 relevés), Țarinii Valley, Bănișoara Sfârș, Ponoare Meadow, Crișului Văratec Valley (1 relevé) (Bihor County), Moneasa Valley (1 relevé) (Arad County). 2 - *Asplenio quadrivalenti-Poëtum nemoralis* association, Brook of Râposu (2 relevés), Tărcăița Valley (5 relevés), Șaua Bălănescu to Summit of Tisa, Moara Dracului-Briheni village, Șopotesei Valley (4 relevés), Rock of Șopotesei, Ponoare Meadow, Șesuța Valley, Summit of Caprei (1 relevé) (Bihor County). 3 - *Asplenietum septentrionali-adianti-nigri* association, Mic Valley (4 relevés), Briheni Valley (5 relevés) (Bihor County). 4 - *Asplenio trichomani-Poëtum nemoralis* association, Șerbanu Valley, Mic Valley, Cusuiș Valley (2 relevés), Briheni Valley, Zărzag Valley (3 relevés), Crișului Văratec Valley, Caselor Hill, Toaca Hill, Tărcăița Valley (Bihor County), Urviș Valley (3 relevés), Archișel Valley, Boroaia Valley (3 relevés), Clit Valley (3 relevés), Zugău Valley, Summit of Izoiu Mic, Răului Valley, Hășmaș Valley (Arad County). 5 - *Ctenidio-Polypodietum* association, Brook of Râposu, Ormanu Valley, Șerban Valley, Morilor Valley - La Stan, Șoim Valley, Moșcoru Valley, Cusuiș Valley, Zărzag Valley, Pont Valley, Finiș Valley, Moara Dracului - Briheni village, Mic Valley, Crișului Văratec Valley, Mare Hill (1 relevé) (Bihor County), Urviș Valley, Clit Valley, Hășmaș Valley, Moneasa Valley, Boroaia Valley, Summit of Izoiu Mic, Summit of Osoiu Mare, Megheș Valley - Piatra Mică, Brook of Osoi, Summit of Merișoara, Răului Valley, Summit of Moma (1 relevé) (Arad County).

In the following table there are shown the values of ecological indices (humidity, temperature, chemical reaction of soil), for the rocky associations studied.

Table 2

The distribution of the species on ecological categories depending on humidity (U), temperature (T), chemical reaction of soil (R)

Asociations	Ecological categories(%)	1	1,5	2	2,5	3	3,5	4	4,5	5	5,5	6	0
As. 1	U	3,12	7,81	28,12	15,63	25	12,5	6,25	-	-	-	-	1,56
	T	-	-	4,69	3,12	50	14,06	15,62	-	1,56	-	-	10,94
	R	-	-	1,56	-	25	-	43,75	-	6,25	-	-	23,44
As. 2	U	-	2,38	7,14	11,91	47,62	21,43	7,14	-	-	-	-	2,38
	T	-	-	9,52	7,14	66,67	2,38	2,38	-	-	-	-	11,91
	R	-	-	-	-	35,71	-	35,71	-	4,76	-	-	23,81
As. 3	U	3,57	-	28,57	14,29	32,14	10,71	7,14	-	-	-	-	3,57
	T	-	-	7,14	7,14	64,29	7,14	7,14	-	-	-	-	7,14
	R	3,57	-	10,71	-	21,43	-	35,71	-	-	-	-	28,57
As. 4	U	1,59	-	7,94	19,05	46,03	17,46	7,94	-	-	-	-	-
	T	-	-	7,94	7,94	68,25	3,17	1,59	-	-	-	-	11,11
	R	-	-	4,76	-	34,92	-	38,1	-	1,59	-	-	20,63
As. 5	U	1,56	4,69	6,25	21,87	37,5	18,75	9,38	-	-	-	-	-
	T	-	-	7,81	6,25	62,5	3,12	6,25	1,56	-	-	-	12,5
	R	-	-	6,25	-	32,81	-	39,06	-	4,69	-	-	17,19

Association: As. 1 - *Asplenietum rutae-murariae-trichomanis*, As. 2 - *Asplenio quadrivalenti-Poëtum nemoralis*, As. 3 - *Asplenietum septentrionali-adianti-nigri*, As. 4 - *Asplenio trichomani-Poëtum nemoralis*, As. 5 - *Ctenidio-Polypodietum*.

Comparative analysis of rocky phytocoenoses behavior from Codru-Moma Mountains is presented in the following ecological spectrum.




Fig. 5 Ecological comparative spectrum, for humidity




Fig. 6 Ecological comparative spectrum, for temperature




Fig. 7 Ecological comparative spectrum, for chemical reaction of soil

CONCLUSIONS

Considering humidity requirements (figure 5), it is observed that for *Asplenio quadrivalenti-Poëtum nemoralis*, *Asplenio trichomani-Poëtum nemoralis* and *Ctenidio-Polypodietum* associations the dominant species are mesophytes ($U_{3-3.5}=56.25-69.05\%$), being the major components of rocky vegetations. The main species for *Asplenietum rutae-murariae-trichomanis* association are the xero-mesophytes ($U_{2-2.5}=43.75\%$), and for *Asplenietum septentrionali-adianti-nigri* association the xero-mesophytes ($U_{2-2.5}=42.86\%$) and mesophytes species ($U_{3-3.5}=42.86\%$) dominate in equal share.

Considering temperature the micro-mesothermophilous species dominate for all the 5 associations studied (figure 6). Thus, on associations, values for the temperature index appears as follows: *Asplenietum rutae-murariae-trichomanis* ($T_{3-3.5}=64.06$), *Asplenio quadrivalenti-Poëtum nemoralis* ($T_{3-3.5}=69.05$), *Asplenietum septentrionali-adianti-nigri* ($T_{3-3.5}=71.43$), *Asplenio trichomani-Poëtum nemoralis* ($T_{3-3.5}=71.43\%$), *Ctenidio-Polypodietum* ($T_{3-3.5}=65.62\%$).

Analysis of species considering the chemical reaction of the soil, highlights the weak acid neutrophylous character for the most part of rocky phytocoenoses identified in the Codru-Moma Mountains (figure 7). The share of weak acid neutrophylous species in the 5 associations studied appears as follows: *Asplenietum rutae-murariae-trichomanis* association ($R_4=43,75\%$), *Asplenio quadrivalenti-Poëtum nemoralis* association ($R_4=35,71\%$), *Asplenietum septentrionali-adianti-nigri* association ($R_4=35,71\%$), *Asplenio trichomani-Poëtum nemoralis* association ($R_4=38,1\%$), *Ctenidio-Polypodietum* association ($R_4=39,06\%$). The large share of basiphile species is explained by the existence of a limestone substrate on extensive areas in Codru-Moma Mountains. Also the presence of the acidophile species is related to the acid substrate, met on high peaks.

REFERENCES

1. Alexiu V., 1998, Vegetația masivului Iezer-Păpușa. Studiu fitocenologic, Edit. Cultura, Pitești, pp. 362.
2. Borhidi A., 2003, Magyarország növénytársulásai, Akadémiai Kiadó, Budapest.
3. Burescu P., Pășcuț C., 2010, Inputs to knowledge of rocks vegetation within Codru-Moma Mountains (NW Romania), „International Conference of Sciences” Analele Universității din Oradea, ISSN 1224-5119, Tom. XVII/1, pp.62-69.
4. Ciocărlan V., 2009, Flora ilustrată a României. *Pteridophyta* et *Spermatophyta*. Editura Ceres, București.
5. Chifu T., Mânzu C., Zamfirescu O., 2006, Flora și vegetația Moldovei (România) vol I-II, Edit. Univ. Al. I. Cuza Iași.

6. Chifu T., Irimia I., Zamfirescu O., 2014, Diversitatea fitosociologică a vegetației României. I Vegetația erbacee naturală. Editura Institutul European, Iași, pp. 603.
7. Cristea V., Gafta D., Pedrotti F., 2004, Fitosociologie. Editura Presa Universitară Clujeană. Cluj-Napoca, pp. 360.
8. Drăgulescu C., 1995, Flora și vegetația din bazinului văii Sadului. Editura Constant, Sibiu, pp. 355.
9. Diaconescu F., 1973, Aspecte din vegetația masivului Leaota, I. Anal. Șt. Univ. "Al. I. Cuza" Iași, Seria nouă, Secț. II-a. Biol. 19, fasc. 2, pp. 465-474.
10. Groza G., 2008, Flora și vegetația Munților Pădurea Craiului. Editura Risoprint, Cluj-Napoca, pp. 300.
11. Karácsonyi C., 2011, Flora și vegetația Dealurilor Tășadului și a colinelor marginale. Rezumatul tezei de doctorat, Arad.
12. Kolbek J., Valachovič M., Mišikova K., 2015, Wall vegetation in old royal mining towns in Central Slovakia. Hacquetia 14/2, pp.249-263.
13. Mihăilescu S., 2001, Flora și vegetația Masivului Piatra Craiului. Edit. Vergiliu București, pp. 400.
14. Niculescu M., 2006, Flora și vegetația bazinului superior al Râului Luncavăț. Rezumatul tezei de doctorat, Cluj-Napoca, pp. 358.
15. Oprea A., Sirbu C., 2009, Munții Stânișoarei (Carpații Orientali). Studiu fitosociologic, Editura Universității „Alexandru Ioan Cuza”, Iași, pp. 214.
16. Pascal P., Mititelu D., 1971, Contribuție la studiul vegetației din bazinul Bistriței Aurii (Jud. Suceava), Comunic. Ști. Inst. Pedagog., Iași, pp.331-363.
17. Paucă A., 1941, Studiu fitosociologic în Munții Codru și Muma. Teză de doctorat M.O., Imprimeria Națională, București, pp. 119.
18. Pășcuț C., 2010, The study of vegetal communities in Codru-Moma Mountains Analele Universității din Oradea, Fascicula: Protecția Mediului, Vol. XV, Anul 15, Editura Universității din Oradea, Oradea, pp. 513-518.
19. Pășcuț C., 2013, Paucă (1941)-Pășcuț (2012) - A comparative study on the vegetation in Codru-Moma Mountains. Analele Universității din Oradea, Fascicula: Protecția Mediului, I.S.S.N. 1224-6255, vol. XX, pp.147-168.
20. Peia P., 1978, Aspecte de vegetație din Cheile Minișului. Contrib. Bot. Cluj-Napoca, pp. 235-250.
21. Pignatti E., Pignatti S., 2014, Plant life of the Dolomites: Vegetation structure and ecology. Springer Publishing House, Rome, pp. 478.
22. Pop I., Cristea V., Hodișan I., 2002, Vegetația județului Cluj. Studiu fitocenologic, ecologic, bioeconomic și ecoprotectiv. Contrib. Bot. (1999-2000), Cluj-Napoca, 35, 2, pp.5-254.
23. Pott R., 1995, Die Pflanzengesellschaften Deutschlands, 2 Aufl., Ulmer Verlag, Stuttgart.
24. Rațiu O., Gergely I., 1976, Structura ecologică și sintaxonomică a vegetației Văii Sebișelului (Bazinul Văii Drăganului). Contrib. Bot., Cluj-Napoca, pp. 73-102.
25. Rațiu O., Gergely I., Șuteu Ș., 1984, Flora și unitățile fitosintaxonomice de pe Valea Iadului (jud. Bihor). Importanța economică și științifică. Caracterizarea lor ecologică III. Contrib. Bot., Cluj-Napoca, pp. 85-135.
26. Răduțoiu D., 2006, Flora și vegetația bazinului Cernei de Olteț. Editura Sintech, Craiova, pp. 407.
27. Sârbu I., Ștefan N., Coroi M., Tănase C., Ciurăscu Ș., 1997, Rezervația naturală Cheile Nărujei - Lacul Negru (jud. Vrancea). Bul. Grăd. Bot., Iași, 6(1), pp. 235-250.

28. Świerkosz K., 2004, Notes on the syntaxonomy of the *Asplenietea trichomanes* class in Poland. Polish Botanical Journal 49(2), pp. 203-213.
29. Ștefan N., Sârbu I., Coroi M., Oprea A., Tănase C., Ciurăscu Ș., 1997, Rezervația naturală Cheile Tișitei (jud. Vrancea). Bul. Grăd. Bot., Iași, 6(1), pp. 251-274.
30. Sanda V., Popescu A., Doltu I., Doniță N., 1983, Caracterizarea ecologică și fitocenologică a speciilor din flora României. Stud. și Com.. Muz. Brukenthal. Sibiu. supliment. 25, pp. 1-126.
31. Sârbu I., Ștefan N., Oprea A., 2013, Plante vasculare din România. Determinator ilustrat de teren. Editura Victor B Victor. București.
32. Schneider-Binder E., 1968, Considerații asupra fitocenozelor de *Asplenium septentrionale* (L.) Hoffm. din Carpații României. Contrib. Bot., Cluj-Napoca, pp. 231-241.
33. Schneider-Binder E., 1969, Contribuții la studiul clasei *Asplenietea rupestris* H. Meier et Br.-Bl. 1934. Contrib. Bot., Cluj-Napoca, pp. 145-155.
34. Schneider-Binder E., 1972, Contribuții la studiul asociațiilor din alianța *Asplenion septentrionalis* Gams 1927 în Carpații României. Muz. Brukenthal, Studii și Comunic. Ști. Nat., Sibiu, 17, pp. 167-181.
35. Stancu I., D., 2005, Flora și vegetația Munților Râiosu și Buda, Masivul Făgăraș. Editura Universității din Pitești.