

STUDIES REGARDING THE INFLUENCE OF CROP ROTATION AND REGIME NUTRITION INTERACTION ON PHYTOMASS ACCUMULATION IN WINTER WHEAT CULTIVATED ON THE PRELUVOSOILS

Bandici Gheorghe*, Ardelean Ileana*, Vidican Teodora*

*University of Oradea, Faculty of Environment Protection, Department of Agricultural, Oradea, Romania
Corresponding author: University of Oradea, Faculty of Environment Protection, Department of Agricultural, 26
Magheru Str., zip code: 410048, Oradea, Romania, tel.: 004059412550, fax. 0040259416274,
e-mail: gbandici@yahoo.com

Abstract

The crop rotation is a decisive factor influencing growth and development of wheat.

Plant growth is fundamental in obtaining yield and is related vegetation and technological factors, the level of yield being reflected in the intensity of phytomass accumulation.

In the majority of cases, total growth of green mass is considered on the assumption that a maximum yield is obtained by increasing total dry weight phytomass production and by a favourable repartition of it among plant's organs.

Bilteanu Gh., (1993) after long tests demonstrated the importance of crop rotation on wheat yields on brown-red soils in Romanian Plain. On clay-illuvial podzols, the introduction of ameliorative plants such as red clover represented an element of outmost importance for increase of the wheat yield.

Key words: crop rotation, regime nutrition, phenophase, phytomass, preluvosoils, early ripening, incomplete ripening, complete ripening

INTRODUCTION

Crop rotation together with other appropriate agricultural practices contribute to the favourableness of growth and development conditions of wheat root system, to an improved synthesis of specific organic compounds and their improved translocation to plant's organs (Lazany, 2000; Bandici ., 1997, 2001) Finally, all the enumerated conditions lead to improved efficiency per area unit.

Zamfirescu, (1977) stressed out that the accumulation process of total green mass relies on the absorption of nutrients from soil through plant's root system. In turn, root system depends on the level of foliar system functioning conditioned by water and nutrients' provision of soil. (Lazany, 2003).

The importance of rationale fertilization and of the crop rotation, on growths and development of plants is stated by many authors: Austin, 1978, Soltner, 1990, Salisbury and Ross, 1995, Domuta et al., 2007, 2008).

Most of the reserches were centred on the influence of crop rotation on the yields, namely on the phytomass accumulation. the crop rotations with regard to wheat was very satisfactory in this order as forerunner plant: pea, beans, winter rape, bots, linseed, soja, red clover, potato, sugar beet, sunflower, corn etc. (Bandici, Gus, 2001, Muntean et al., 2011, Domuta, 2012).

Dincă, (1982), made some references on the role of crop rotation on wheat yield and on the organic accumulation in whole plant and grains.

It is demonstrated that after 10-year monoculture, wheat yield decreases continously in comparative with rotations. it fluctuates as a consequence of changing climatic conditions. under such circumstances, fertilization does not induce a significant yield increase. a particularly important problem is linked to wheat crop increment, which must fit the rising consumption needs of world population (Zăhan 1989, Bandici et al., 2003, Ardelean, 2013).

MATERIAL AND METHOD

The research was set at Agricultural Researche and Development Station (A.R.D.S) Oradea (Romania), between 2018-2019. The experimental design was polyfactorial in subdivides stands using as factors interaction: crop rotation x phenophase and regime nutrition x phenophase.

The influence of later factors was studied on the dynamics phytomass accumulation of winter wheat cultivated on preluvosols. As biological material, the Delia race of wheat was employed.

Experimental results (phytomass accumulation) were analyzed by ANOVA (analysis of variance) and expressed as g of dry weight/10 plants (Bandici, 2007).

RESULTS AND DISCUSSION

The present paper presents the correlation between phytomass accumulation and crop rotation quality in winter wheat cultivation.

The Table 1 presents the phytomass accumulation during the less favourable year 2018. The results show that the forerunner plant do not influence phytomass accumulation increase as winter wheat advances in vegetation. The increase is dependent on the properties and quality of the crop rotation, the best yields being obtained after the cultivation of pea.

Thus, in wheat monoculture the phytomass accumulation increases from 0.96 g. dry weight/10 plants (at beginning winter) to 45.40 g. dry weight/10 plants at complete ripening. If corn is used as crop rotation, but specially after pea, the increases in yield are superior, varying between 0.87-

60.43 g. dry weight/10 plants after corn and between 1.50-66.97 g. dry weight/10 plants after pea, respectively as winter wheats' forerunner plants.

The phenomenon keeps same pace but at values in 2019, a less favourable year (table 2). The phytomass accumulation varies in wheat monoculture between 0.20-35.60 g. dry weight/10 plants and 0.23-42.83 g. dry weight/10 plants after corn and between 0.20-47.10 g. dry weight/10 plants after pea. Regardless to climatic conditions, at a different scale, a positive correlation was found between phenophase x regime nutrition: as wheat advances to maturity, there is a progressive accumulation of total phytomass in seeds (table 3).

Concerning 2018, results show a very significant increase in phytomass at the beginning of winter in unfertilized alternative (1.00 g dry weight/10 plants). At the complete ripening there is an increase to 49.92 g dry weight/10 plants, in unfertilized alternative (Zăhan, 1989).

In 2019, a less favorable year, phytomass accumulation decreases as compared to a favorable year, taking values in a narrower, range between 0.22 g dry weight/10 plants at beginning of winter and 31.32 g dry weight/10 plants at compared ripening in unfertilized alternative.

It is worth to mention that in 2018 the quantity of accumulated phytomass was directly proportional with fertilization level as it was rising during study period (49.92 g dry weight/10 plants) in unfertilized alternative and 69.49 g dry weight/10 plants in fertilized alternative, organo-mineral complex was used in all experimental alternative as fertilized.

In a less favorable year, 2019, phytomass accumulation decreased depending on nutrition regime (31.32 g dry weight/10 plants in unfertilized alternative and 51.80 g dry weight/10 plants in fertilized alternative using the same organo-mineral complex as fertilizer).

It is worth to mention that during the two research years, a negative correlation was found between percent participation of compound synthesized before fructification and seeds formation and fertilization level.

This phenomenon is more accentuated in unfertilized alternatives as compared with fertilized with organo-mineral complex and is influenced unfavorable climatic conditions.

Table 1

The effect of nutrition regime x phenophase interaction on winter wheat dry weight phytomass accumulation dynamics on preluvosols.
(Oradea 2018)

Phenophase	Total dry weight phytomass, seeds and straw g./10 plants								
	Nutrition regime								
	N ₀ P ₀			N ₁₂₀ P ₈₀			N ₁₀₀ P ₈₀ + 10t/ha manure		
	Total s d.W.	Seeds	Straw	Total s d.W.	Seeds	Straw	Total s d.W.	Seeds	Straw
At winter beginning	1.00	-	1.00	1.17	-	1.17	0.97	-	0.97
At the end of winter	1.22	-	1.22	1.55	-	1.55	1.65	-	1.65
The beginning of vegetation	2.65	-	2.65	2.62	-	2.62	4.65	-	4.65
The formation of first interned	4.77	-	4.77	6.10	-	6.10	7.45	-	7.45
Straw elongation	11.62	-	11.62	16.85	-	16.85	19.37	-	19.37
The formation of spike	36.70	-	36.70	43.05	-	43.05	44.10	-	44.10
Beginning of seeds formation	43.62	-	43.62	49.55	-	49.55	49.31	-	49.31
Early ripening	47.62	9.50	38.12	52.90	13.75	39.15	56.45	13.25	43.20
Incomplete ripening	49.82	15.75	34.07	57.32	18.45	38.87	63.55	17.40	46.15
Complete ripening	49.92	21.00	28.92	57.95	24.50	33.45	69.49	27.75	41.74
LSD 5%	0.063 g/10 plants d.w.						2.3 g/10 plants d.w.		
LSD 1 %	For total plant phytomass: 0.083 g/10 plants d.w.						For seeds: 3.1 g/10 plants d.w.		
LSD 0.1 %	0.106 g/10 plants d.w.						4.0 g/10 plants d.w.		

Statistical significations:

- for *Total plant phytomass*: under 0.063 = insignificant (-); 0.063-0.083 = significant (*); 0.083-0.106 = distinct significant (**); over 0.106 = very significant (***)

- for *seeds*: under 2.3 = insignificant (-); 2.3-3.1 = significant (*); 3.1-4.0 = distinct significant (**); over 4.0 = very significant (***)

Table 2.

The effect of the crop rotation x phenophase on winter wheat phytomass accumulation dynamics. on preluvosols (Oradea, 2019)

Phenophase	Total dry weight phytomass. seeds and straw g./10 plants											
	Crop rotation											
	Wheat monoculture (Mt)			Corn (W-C)			Pea (P-W-C)			Pea (P-W-C-C)		
	Total s d.W.	Seeds	Straw	Total s d.W.	Seeds	Straw	Total s d.W.	Seeds	Straw	Total s d.W.	Seeds	Straw
At winter beginning	0.20	-	0.20	0.23	-	0.23	0.30	-	0.30	0.20	-	0.20
At the end of winter	0.37	-	0.37	0.37	-	0.37	0.40	-	0.40	0.37	-	0.37
The beginning of vegetation	0.47	-	0.47	0.47	-	0.47	0.50	-	0.50	0.47	-	0.47
The formation of first interned	0.80	-	0.80	0.97	-	0.97	1.33	-	1.33	0.93	-	0.93
Straw elongation	3.73	-	3.73	3.43	-	3.43	4.37	-	4.37	4.33	-	4.33
The formation of spike	8.60	-	8.60	9.47	-	9.47	11.0	-	11.00	10.60	-	10.60
Beginning of seeds formation	24.83	-	24.83	25.50	-	25.50	34.03	-	34.03	33.13	-	33.13
Early ripening	33.87	9.83	24.04	36.00	8.50	27.50	35.73	11.33	24.40	41.33	10.83	30.50
Incomplete ripening	35.37	11.70	23.67	41.87	11.00	30.87	40.73	13.90	26.83	45.33	12.17	33.16
Complete ripening	35.60	13.37	22.23	42.83	12.80	30.03	43.43	17.50	25.93	47.10	15.97	31.13
LSD 5%	0.073 g/10 plants d.w.						2.7 g/10 plants d.w.					
LSD 1 %	.For total plant phytomass: 0.096 g/10 plants d.w.						For seeds: 3.6 g/10 plants d.w.					
LSD 0.1 %	0.123 g/10 plants d.w.						4.7 g/10 plants d.w.					

Statistical significations:

- for *Total plant phytomass*: under 0.073 = insignificant (-); 0.073-0.096 = significant (*); 0.096-0.123 = distinct significant (**); over 0.123 = very significant (***)

- for *seeds*: under 2.7 = insignificant (-); 2.7-3.6 = significant (*); 3.6-4.7 = distinct significant (**); over 4.7 = very significant (***)

Table 3

The effect of nutrition regime x phenophase interaction on winter wheat dry weight phytomass accumulation dynamics on preluvo soils
(Oradea 2019)

Phenophase	Total dry weight phytomass. seeds and straw g./10 plants								
	Nutrition regime								
	N ₀ P ₀			N ₁₂₀ P ₈₀			N ₁₀₀ P ₈₀ + 10t/ha manure		
	Total s d.W.	Seeds	Straw	Total s d.W.	Seeds	Straw	Total s d.W.	Seeds	Straw
At winter beginning	0.22	-	0.22	0.22	-	0.22	0.25	-	0.25
At the end of winter	0.35	-	0.35	0.37	-	0.37	0.40	-	0.40
The beginning of vegetation	0.45	-	0.45	0.47	-	0.47	0.50	-	0.50
The formation of first interned	0.90	-	0.90	0.95	-	0.95	1.17	-	1.17
Straw elongation	2.45	-	2.45	4.47	-	4.47	4.97	-	4.97
The formation of spike	5.82	-	5.82	11.30	-	11.30	12.62	-	12.62
Beginning of seeds formation	23.00	-	23.00	30.07	-	30.07	35.05	-	35.05
Early ripening	26.45	6.03	20.42	38.97	11.85	27.12	44.77	12.50	32.27
Incomplete ripening	30.22	7.35	22.87	43.05	14.07	28.98	49.20	15.15	34.05
Complete ripening	31.32	9.72	21.60	43.60	15.70	27.90	51.80	19.30	32.50
LSD 5%				0.063 g/10 plants d.w.			2.3 g/10 plants d.w.		
LSD 1 %	For total plant phytomass:			0.083 g/10 plants d.w.			For seeds: 3.1 g/10 plants d.w.		
LSD 0.1 %				0.106 g/10 plants d.w.			4.0 g/10 plants d.w.		

Statistical significations:

- for *Total plant biomass*: under 0.063 = insignificant (-); 0.063-0.083 = significant (*); 0.083-0.106 = distinct significant (**); over 0.106 = very significant (***)

- for *seeds*: under 2.3 = insignificant (-); 2.3-3.1 = significant (*); 3.1-4.0 = distinct significant (**); over 4.0 = very significant (***)

Thus, in 2018, a favorable year, increasing the fertilization had as result a decrease of participation percentage from 33.7 g dry weight/10 plants in unfertilized alternative to 15.3 g dry weight/10 plants in fertilized with organo-mineral complex alternatives.

In 2019, considered an unfavorable year, the decreases varied between 11.2 g dry weight/10 plants in unfertilized and 7.9 g dry weight/10 plants in fertilized alternatives using organo-mineral complex of fertilizers.

CONCLUSIONS

The crop rotation has a positive effect on total phytomass accumulation as compared with wheat monoculture, the obtained values being conditioned by a higher favourableness of climatic factors.

With regard to phytomass accumulation in seeds, it was positively influenced by the quality of the crop rotation, being higher in correlation with the nutrition regime qualities.

REFERENCES

1. Ardelean I., 2013, Agrotehnica. Editura Universității din Oradea. p.417
2. Austin R.B., 1978, ADAS, Qualerly Review, 29, 76-87.12
3. Bandici G. E., 1997, Contributii la stabilirea influenței premergătoarei si a fertilizării asupra dinamicii acumularii biomasei, la grâul de toamna, cultivat pe soluri cu exces temporar de umiditate, în centrul Câmpiei de Vest a României. Doctoral thesis. University of Agriculture Sciences and Veterinary Medicine Cluj-Napoca, Romania, p.250.
4. Bandici G. E., C. Domuta, I. Ardelean, 2003, The influence of the forerunner plant, fertilisation level and climatic conditions on the total wet and dry gluten content of winter wheat seeds cultivated on brown luvic soils in the Western Plain of Romania. *Lucrari știintifice USAMVB.*, Seria B, vol. XLV, Bucuresti, p.281-284, p.330.
5. Bandici, G.E., Guș, P., 2001, Dinamica acumulării de biomasă la grâul de toamnă. University of Oradea Press, 107 p.
6. Bîlteanu, G., 1993, Fitotehnie, Ceres Printing House. Bucharest, pp. 457.
7. Dinca D., 1971, Influenta rotatiei asupra productiei, valorificarii îngrasamintelor si calitatii biologice a recoltelor de grâu si porumb pe solul brun roscat de padure. *Probleme agricole*, no.9, p.56-59, p.70.
8. Dincă, D., 1982, Asolamentele agriculturii moderne. Ceres Printing House. Bucharest. 257 pp
9. Domuta C., 2012, Agrotehnică. Editura Universității din Oradea. p.506
10. Domuta C., Bandici Gh., Ciobanu Gh., N. Csep, Ciobanu C., Samuel A., Bucurean E., Sandor M., Borza I., Bunta Gh., Ileana Ardelean, Cr. Domuta, 2007, "Asolamentele în Câmpia Crisurilor". Editura Universitatii din Oradea, ISBN 978-973-759-350-4, pag. 254.
11. Domuta C., Bandici Gh., Ciobanu Gh. Ciobanu C., Samuel A., N. Csep, Bucurean E., Borza I., Sandor M., Bunta Gh., I. Ardelean, Cr. Domuta., 2008, Asolamentele in sistemele de agricultura, Editura Universitatii din Oradea, pag. 297..
12. Lazany, J., 2000, Soil fertility management in Westik's crop rotation experiment. Role of fertilizers in Sustainable Agriculture. CIEC Conference. pp.77-80.
13. Lazany, J., 2003, Differences in soil carbon content in the treatments of Westik's crop rotation experiment. Natural resources and sustainable development. International scientific session and reviewed papers. Oradea-Debrecen, pp. 119-120.

14. Muntean L. S., S. Cernea, G. Morar, M. Duda, D. Vârban, S. Muntean, 2011, Fitotehnie. Academic Pres Printing House, Cluj-Napoca, p.83-135, p.255.
15. Salisbury F.B., C.W. Ross, 1995, Fisiologia vegetale. Seconda edizione italiana condota sulla quarta edizione americana. Editura Zanichelli..
16. Soltner D., 1990, Phytotechnie speciale, Colection sciences et Techniques Agricoles, Angers.
17. Zamfirescu, N., 1977, Bazele biologice ale producției vegetale. Ceres Printing House, Bucharest, 337 p.
18. Zăhan, P., Zăhan, R., 1989, Cercetări privind influența plantei premergătoare și a fertilizării asupra dinamicii de acumulare a masei vegetale la grâul cultivat pe soluri podzolice cu exces temporar de umiditate din Câmpia de Vest a țării (I). Probleme de agrofitotehnie teoretică și aplicată nr. 1, vol. XI: 97-102.