

DOSES AND EFFECTS INCREASE THE AMOUNT OF UV-B RADIATION ON PLANT *ZEA MAYS.L.*

Orodan Maria^{*}, Morgovan Claudiu^{*}, Olah Neli-Kinga^{*}, Atyim Paul^{*}, Negrei Carolina^{}, Toderescu Corina Dalia^{*}, Ginghina Octav^{**}, Hulbar Aurelia^{*}, Borza Corina^{*}, Osser Gyongyi^{*}**

^{*}”Vasile Goldis” Western University of Arad, Faculty Pharmacy 91-93, L. Rebreanu Street, Arad, Romania

^{**}“Carol Davila” University of Medicine and Pharmacy, Department of Toxicology 6, Traian Vuia Street, Bucharest, Romania
e-mail: gyongyiosser@ymail.com

Abstract

The utilized radiation dose and the action time varies, depending on the tested species, but the simplicity of the required gear and the in vitro obtained effects make UV irradiation the main method of inducing mutations

The UV-B waves are not only a stress factor, but, as part of the Sun’s electromagnetic wave emissions, have useful information about the waves’ relations with the environmental factors (the ozone layer, the irradiation angle, topographic factors and leaf development).

Following these changes, the effect may be lethal if the reparatory process is not carried out. To counteract the negative UV radiation effects, the organisms and plants have regenerative mechanisms (photoreactivation, excision repair, recombination replicative repair).

Key words: *Zea mays L.*, UVB radiation, reparatory process

INTRODUCTION

The UV radiation can cause chemical bonds to break and photochemical reactions.

The effects of the UV radiation on plants (particularly UV-B ones) vary, depending on the species and on the variety of said species.

Although with the increase of the UV-B radiation intensity, the growth and development of superior plants is usually negatively affected, the magnitude of the effect depends on the taxonomic position of the plant, on the growth culture medium conditions and on its ontogenetic stage of development (Rao et al, 1996, Rozema et al, 2002, Roberts, 1965).

Of all the studied plant species, between 30% and 50% are negatively affected by the increase in the amount of UV-B radiation.

The remainder of the tested species were either not affected or seemed to benefit from the increased UV-B radiation levels (Caldwell et al, 1989, Cachiță, et al, 2004).

The effects of the increased UV radiation amount also include the reduction of the biomass, photosynthesis inhibition, the induction of

secondary vegetal compounds, photomorphogenic and phonological changes (Sancar, 1994, Pallag et al. 2014, Pop et al, 2011, Perry et al, 1999, Casati &. Walbot, 2004, Caldwell et al, 1989).

The most susceptible nucleotides to the action of the UV radiation are the ones containing pyrimidine bases (thymine and cytosine).

The effect characteristic to UV irradiation is the forming of pyrimidine dimers, which distorts the shape of the DNA molecule, perturbing the replication (Pallag et al. 2014, Perry et al, 1999, Casati &. Walbot, 2004, Cachiță et al, 2004, Bungau et al, 2011, Basiouny, 1986, Bungău, 2014, Băra et al, 2003).

MATERIAL AND METHOD

Many of the agricultural plants of subtropical in origin (for example, corn, rice, tomatoes, peppers, squash, cucumbers etc.) are also found in the temperate zone, but are cultivated under less than optimal conditions.

In this case, the reproduction of these plants must have the role of selecting the genotypes with the appropriate tolerance to the imposed conditions and, via them, creating varieties that are enduring (Băra & Crețu, 2006, Băra,et al, 2003, Bungău et al, 2011, Bungău et al, 2003).

Under the effect of the treatment with UV-B wavelength, the increase in the POD activity, by comparing the control batch and the treated one, for the Helga hybrid maize plants, was insignificant after a day of irradiation; it became significant after 4 days of irradiation.

For the 287 nm wavelength, on the 1st, 4th and 7th days, samples were taken from the third, fully developed leaf, for the analysis of the antioxidant enzymes activity: CAT, and POD, in accordance with the methods described in the specialty literature.

The increased ROS levels determine the significant increase of the antioxidant enzymes' activity, a process by which the plants are protected against the damaging effects of the oxidative stress.

The increase in the activity of the antioxidant defensive system, due to the negative effects of the environmental stressors, is developed over a long period of time; this process is called adaptation (Băra & Crețu, 2006, Băra,et al, 2003, Bungău et al, 2011, Bungău et al, 2003, Caldwell et al, 1989, Rozema et al, 2002).

RESULTS AND DISCUSSIONS

Based on the available data from the literature, we asked ourselves if the number of days under the influence of radiation and UV-B induce changes dependent on the spectrum used on the antioxidant system in plants.




Fig.1 The change in absorbance under the effect of the UV-B wavelengths, regarding the activity of the POD enzyme for the HELGA and ZP471 hybrids

In the case of CAT, on the 4th day of treatment, one could notice a decreasing tendency in its accumulation in the leaves, whilst GR and GST remained active until the last day of treatment (Fig. 1).

The APX activity increased to a great extent, due to activation under the wavelength action, reaching its peak in the 4th day. In the first day, the POD activity did not show any changes, whilst in the 4th day it only increased slightly, but not in a significant way (Fig. 2).




Fig.2. The change in absorbance under the effect of the UV-B wavelengths, regarding the activity of the APX enzyme for the HELGA and ZP471 hybrids

CONCLUSIONS

In their natural environment, the plants are subjected to stress factors, including UV-B radiations (280 – 320 nm) which play an important role, as an increasing number of UV-B radiations reaches Earth's surface because of the stratospheric ozone layer's depletion.

Following the treatment we noticed differences in the activation of certain enzymes, regarding the accumulation degree and rhythm.

Following the treatment, the GR, GST and CAT enzymes were similarly activated for the control plants as well as the treated ones, in the first day; however, in the 4th day, the UV-B effect determined differentiations between the control batch and the treated one, where statistically significant values were recorded.

For the 7th day, all three enzymes decreased their activity, with more significant differences in the case of CAT.

REFERENCES

1. Basiouny, F.M., 1986, Sensitivity of corn, oats, peanuts, rice, rye, sorghum, soybean and tobacco to UV-B radiation under growth chamber conditions, *J. Agron. Crop Sci*, vol 157:31–35.
2. Băra C. I., Crețu R. M., 2006, Biochemical effects induced by UV treatment on 5 romanian *Phaseolus vulgaris* L. cultivars, grown in field” *Scientific Annals of Universității “Al.I.Cuza” University of Iași (new series), Section I, a. Genetics and Molecular Biology*, vol 8:145-150.
3. Băra, C., Artenie, V, Băra, I. I., 2003, Effects of the UV-B radiations in superior plants” *Scientific Annals of Universității “Al.I.Cuza” University of Iași (new series), Section I, a. Genetics and Molecular Biology*
4. Bungău, S., 2014, Vitamins and amino acids determination using kinetic methods, *Colecția Science&Technology*, Ed. Italian Academic Publishing, București
5. Bungău S., Fodor, A., Szabo, I., Muțiu, G., 2011, Comparative studies on *Petroselinum crispum folium* ascorbic acid content using kinetic, spectrophotometric and iodometric methods, *Archives of the Balkan Medical Union*, 46(1), pp. 77-80
6. Bungău, S., Fodor, A., Țiț, D. M., Szabó, I., 2011, Studies on citrus species fruits ascorbic acid content using kinetic, spectrophotometric and iodometric methods, *Analele Universității din Oradea, Fascicula Protecția Mediului*, vol. 16(A):212-217
7. Bungău, S., Bâldea, I., Copolovici, L., 2003, Determination of ascorbic acid in fruit using a Landolt type method, *Revista de Chimie*, 54 (3), pp. 213-216
8. Cachiță, C.D., Petruș, C.M., Petruș-Vancea, A., Crăciun, C., 2008. „Hyperhydricity phenomenon developed at the level of sugar beet (*Beta vulgaris* L. Var. *Saccharifera*) vitrocultures, III. Ultrastructural aspects observed in the hyperhydric callus cells. *Studia Univ. „V. Goldis”*, 18(Suppl):53-64.
9. Caldwell, M. M., Robberecht, R., Flint, S. D., 1983, „Internal filters: Prospects for UV-acclimation in higher plants.” *Physiol. Plant.* 58, Copenhagen, p. 445-450.
10. Caldwell, M. M., Teramura, A. H., Tevini, M., 1989. The changing solar ultraviolet climate and the ecological consequences for higher plants, *Trends in Ecology and Evolution*, (4):363-367.
11. Casati, P. and Walbot, V., 2004. „Rapid transcriptome responses of maize (*Zea mais*) to UV-B, in irradiated and shielded tissues.” *Genome* 5, R 16
12. Paul ND, Gwynn-Jones D., 2003, Ecological roles of solar UV radiation: towards an integrated approach”. *Trends Ecol Evol*, vol 18:48–55.
13. Perry NB, Anderson RE, Brennan NJ, Douglas MH, Heaney AJ, McGimpsey JA, Smallfield BM ,1999. „Essential oils from Dalmatian (*Salvia Officinalis* L.): variations among individuals, plant parts, seasons, and sites”. *J. Agric. Food Chem.*, 47, p. 204-205.
14. Pop, T.I., D. Pamfil, C. Bellini, 2011, Auxin control in the formation of adventitious roots, *Notulae Botanicae Horti Agrobotanici Cluj-Napoca*, vol 39(1):307-316.
15. Pallag A., Osser Gy., Orodan M., Gîtea D., Bungău S., Pașca B., 2014, Palynologic Analysis Of Allergenic Species In Oradea and Arad (Romania) Area Between 2012-2014, *Analele Universității din Oradea, Fascicula Protecția Mediului*, vol 23:867-872.
16. Pallag A., Pașca B., Gîtea D., Bungău S., Țiț D. M., 2014, The Influence Of Some Microelements on Germination and Chlorophyll Quantity In *Ocimum basilicum* L. *Analele Universității din Oradea, Fascicula Protecția Mediului*, vol 23:873-880.
17. Pallag A., Pașca B., Gîtea D., Țiț M.D., 2014, The effects of acetylsalicylic acid in physiological processes of *Triticum aestivum* L. *Analele Universității din Oradea, Fascicula: Fascicula Protecția Mediului*, Vol. 23:119-124.

18. Pallag A., Osser Gy., Orodan M., Honiges A., Gîtea D., Paşca B., 2014, Determining the amount of nucleic acids in medicinal ferns collected from different areas of ARAD and BIHOR county, *Analele Universităţii din Oradea, Fascicula Protecţia Mediului*, vol 23:263-268.
19. Rao M.V., Paliyath, G., Ormrod, D.P., 1996. Ultraviolet-B and ozone-induced biochemical changes in antioxidant enzymes of *Arabidopsis thaliana*". *Plant Physiology* (Rockville), vol 110 (1):125-136.
20. Roberts, L.G., 1965, Machine perception of three-dimensional solids. *Optical and Electro-Optical Information Processing*, 159-197.
21. Rowland R.A., 1991, Physiological and morphological responses of snapbean plants to ozone stress as influenced by pretreatment with UV-B radiation". In: Oxford & IBH Publishing Co.Pvt. Ltd., New Delhi, p. 133–146.
22. Rozema J, Geel B, Bjorn LO, Lean J, Madronich S., 2002, Toward solving the UV puzzle, *Science*, vol 296:162–162.
23. Sancar A., 1994, Structure and function of DNA photolyase, *Biochemistry* vol 33:2-9.